arXiv:0806.4939v1 [math.QA] 30 Jun 2008

MATRIX FACTORIZATIONS AND INTERTWINERS OF THE FUNDAMENTAL REPRESENTATIONS OF QUANTUM GROUP $U_q(\mathfrak{sl}_n)$

YASUYOSHI YONEZAWA

Abstract. We want to construct a homological link invariant whose Euler characteristic is MOY polynomial as Khovanov and Rozansky constructed a categorification of HOMFLY polynomial. The present paper gives the first step to construct a categorification of MOY polynomial. For the essential colored planar diagrams with additional data which is a sequence naturally induced by coloring, we define matrix factorizations, and then we define a matrix factorization for planar diagram obtained by gluing the essential planar diagrams as tensor product of the matrix factorizations for the essential planar diagrams. Moreover, we show that some matrix factorizations derived from tensor product of the essential matrix factorizations have homotopy equivalences corresponding to MOY relations.

Contents

1. Introduction 1
1.1. Categorification of quantum link invariant 1
2. Category of $\mathbb{Z}$-graded matrix factorization 5
2.1. $\mathbb{Z}$-graded matrix factorization 5
2.2. Koszul matrix factorization 7
3. Matrix factorizations for colored planar diagrams 13
3.1. Poincaré series of a $\mathbb{Z}$-graded algebra 13
3.2. Colored planar diagrams and matrix factorizations 14
References 36

1. Introduction

1.1. **Categorification of quantum link invariant.** Mikhail Khovanov introduced a homological link invariant whose Euler characteristic is Jones polynomial. At present we understand Jones polynomial as a link invariant derived from the quantum group $U_q(\mathfrak{sl}_2)$ and its 2-dimensional vector representation V_2. (We can also obtain a link invariant induced by a quantum group and its representation, called a quantum link invariant.) Such a system making a homological link invariant whose Euler characteristic is a quantum link invariant is called a categorification of the quantum link invariant. A natural question is "can we construct a categorification of the other quantum link invariants?" In the case of HOMFLY polynomial, which is derived from $U_q(\mathfrak{sl}_n)$ and its n-dimensional representation V_n, Mikhail Khovanov and Lev Rozansky also constructed a homological link invariant whose Euler characteristic is HOMFLY polynomial.



Figure 1. Reductions for single crossings of HOMFLY polynomial

---

2000 *Mathematics Subject Classification.* 81R50,18G60.

Key words and phrases. matrix factorization, categorification, Khovanov-Rozansky homology.

However, there exist a lot of quantum link invariants which is not categorified yet. For example, MOY polynomial (see [8]), which is a quantum (regular) link invariant derived from $U_q(\mathfrak{sl}_n)$ and its fundamental representations, is one of quantum link invariants which are not categorified (uncategorified) yet. Since this MOY polynomial is a generalization of HOMFLY polynomial, it is natural that we hope to construct a categorification of MOY polynomial generalizing Khovanov and Rozansky's work.

We briefly recall the work of M.Khovanov and L.Rozansky [5]. We calculate HOMFLY polynomial of an oriented link diagram D by transforming each single crossing into planar diagrams P_0 and P_1 in Figure 2 (in the case of Hopf link, see Figure 3), evaluating the planar diagrams Γ derived from the link diagram as a Laurent polynomial of q and summing the Laurent polynomials by the reduction in Figure 1.

For an oriented link diagram D, we can construct a homological link invariant $\mathcal{C}(D)_n$ whose Euler characteristic is HOMFLY polynomial for the link diagram D as follows. First, for each planar diagram P in Figure 2 we define a matrix factorization $\mathcal{C}(P)_n$ which is a 2-cyclic complex of modules. Since every planar diagram Γ induced by a link diagram D is decomposed into some of the planar diagrams P_0 and P_1, a matrix factorization $\mathcal{C}(\Gamma)_n$ for the planar diagrams Γ is defined to be the tensor product of matrix factorizations for parts P_0 and P_1 of the decomposition. For example, the planar diagram Γ_{10} in Figure 3 has a decomposition shown as Figure 4. Then we have a matrix factorization $\mathcal{C}(\Gamma_{10})_n$ for the diagram Γ_{10} as



Figure 2. Planar diagrams



Figure 3. Planar diagrams derived from Hopf link diagram

$$\mathcal{C}(\Gamma_{10})_n = \mathcal{C}(\Gamma_1)_n \boxtimes \mathcal{C}(\Gamma_2)_n \boxtimes \mathcal{C}(\Gamma_3)_n.$$



Figure 4. Decomposition of planar diagram

For a link diagram D, we define a complex $\mathcal{C}(D)_n$ by exchanging every crossing into a complex of matrix factorizations for planar diagrams P_0 and P_1 as follows,



In the case of Hopf link diagram, we obtain the following complex,

$$\mathcal{C}(\Gamma_{00})_n \xrightarrow{\begin{pmatrix}\chi_-\boxtimes\mathrm{Id}\\ \mathrm{Id}\boxtimes\chi_-\end{pmatrix}} \mathcal{C}(\Gamma_{10})_n\oplus\mathcal{C}(\Gamma_{01})_n \xrightarrow{(\mathrm{Id}\boxtimes\chi_-,-\chi_-\boxtimes\mathrm{Id})} \mathcal{C}(\Gamma_{11})_n \ .$$

M. Khovanov and L. Rozansky introduced such a complex of matrix factorizations for a link diagram D and they proved that χ_+ and χ_- of the complex are associated with a homological link invariant, i.e. they showed that complexes derived from diagrams appearing in Reidemeister moves are isomorphic.

We want to construct a homological link invariant whose Euler characteristic is MOY polynomial in [8] as Khovanov and Rozansky constructed a categorification of HOMFLY polynomial. The present paper gives the first step to construct a categorification of MOY polynomial. We consider the following strategy to achieve this purpose,

Strategy 1.1 (Categorification of MOY polynomial)**.**

(S1) *Give a graded category* MF^{gr} *of a* $\mathbb{Z}$*-graded matrix factorization with tensor product (commutativity and associativity) such that some matrix factorization* $\mathcal{C}(P)$ *of* MF^{gr} *realizes every essential colored planar diagram* P *of Figure* 5*.*



$$1\le i\le n, 1\le i_1,i_2\le n-1, i_3=i_1+i_2\le n$$

Figure 5. Essential colored planar diagrams

By calculation of MOY polynomial (see [8]*) every colored crossing is exchanged into a formal linear sum of* Γ_k^L *and* Γ_k^R *in Figure* 6*. Since the diagrams* Γ_k^L *and* Γ_k^R *is decomposed into some of essential planar diagrams in Figure* 5*, every colored planar diagram* Γ *derived from a colored link diagram* D *is also decomposed into some of essential planar diagrams in Figure* 5*. For the colored planar diagram* Γ*, we define a matrix factorization* $\mathcal{C}(\Gamma)$ *to be tensor product of matrix factorizations for parts of the decomposition.*

Moreover, the category MF^{gr} *must have a homotopy equivalence (i.e. an isomorphism in the homotopy*



Figure 6. Colored planar diagrams

category HMF^{gr}*) corresponding to every relation of MOY polynomial. For example, the homotopy category* HMF^{gr} *must have the following equivalence corresponding to MOY relation in* [8]*,*



(S2) *In such a homotopy category* HMF^{gr}*, we construct* $\mathbb{Z}$*-grading preserving morphisms* $\chi_{+,k}^{(i,j)}$ *from* $\mathcal{C}(\Gamma_k^R)$ *to* $\mathcal{C}(\Gamma_{k-1}^R)$ *and morphisms* $\chi_{-,k}^{(i,j)}$ *from* $\mathcal{C}(\Gamma_k^L)$ *to* $\mathcal{C}(\Gamma_{k+1}^L)$*. Using these morphisms* $\chi_{+,k}^{(i,j)}$ *and* $\chi_{-,l}^{(i,j)}$*, we define the chain complex for colored single crossings as a complex of matrix factorizations in the complex category* $\mathrm{Kom}(\mathrm{HMF}^{gr})$*. Then we prove that such morphisms* $\chi_{+}^{(i,j)}$ *and* $\chi_{-}^{(i,j)}$ *give a homological link invariant. That is, we have to prove that the complexes defined by these morphisms* $\chi_{+,k}^{(i,j)}$ *and* $\chi_{-,k}^{(i,j)}$ *induce the following isomorphism in the homotopy category of* $\mathrm{Kom}(\mathrm{HMF}^{gr})$



The main purpose of this paper is defining a candidate for matrix factorizations in Strategy 1.1 (S1). For the essential colored planar diagrams (Figure 5) with additional data which is a sequence naturally induced by coloring, we define matrix factorizations, and then we define matrix factorizations for planar diagrams Γ_k^R and Γ_k^L of Figure 6 as tensor product of matrix factorizations for essential planar diagrams. In the paper [10], the author defined matrix factorizations for the planar diagrams ,  and  to reconstruct the matrix factorization for . Since the planar diagrams ,  and  are the same to colored planar diagrams ,  and , we can define matrix factorizations for diagrams of Figure 5 as a generalization of these matrix factorizations.

In the notion of the category HMF^{gr}, we show that some matrix factorizations derived from tensor product of the essential matrix factorizations have isomorphisms in HMF^{gr} corresponding to MOY relations.

The present paper is organized as follows.

In Section 2, we give basic definitions, properties and theorems related to category of a matrix factorization. However, many things are already known. New results are Theorem 2.13, which is a generalization to multivariable of Theorem 2 given by Khovanov and Rozansky [7][10], Corollary 2.11, Corollary 2.14 and Corollary 2.15. In Section 3, we notice facts of $\mathbb{Z}$-graded algebra. After that, we define matrix factorizations for essential colored planar diagrams of Figure 5 and show that some matrix factorizations for planar diagrams have equivalences corresponding to MOY relation.

Acknowledgement. The author is grateful to O. Iyama, H. Ochiai and K. Yoshida, Y. Yoshino, especially A. Tsuchiya and L. Rozansky for their helpful comments. This work is partly supported by the Grant-in-Aid for JSPS Fellows (20-2330) from Japan Society for the Promotion of Science.

2. Category of $\mathbb{Z}$-graded matrix factorization

2.1. **$\mathbb{Z}$-graded matrix factorization.** Let R be a $\mathbb{Z}$-graded polynomial ring over $\mathbb{Q}$, let M_0,M_1 be free $\mathbb{Z}$-graded R-modules permitted infinite rank and let be a homogeneous element $\omega \in R$. A **matrix factorization** with a potential $\omega \in R$ is a 2-cyclic complex of M_0 and M_1,

$$\overline{M} = \left(M_0 \xrightarrow{\quad d_{M_0} \quad} M_1 \xrightarrow{\quad d_{M_1} \quad} M_0 \right)$$

with the properties $d_{M_1} d_{M_0} = \omega \,\mathrm{Id}_{M_0}$ and $d_{M_0} d_{M_1} = \omega \,\mathrm{Id}_{M_1}$. We often simply denote this 2-cyclic complex by $\overline{M} = (M_0, M_1, d_{M_0}, d_{M_1})$.

For a $\mathbb{Z}$-graded polynomial ring R over $\mathbb{Q}$ and $\omega \in R$, let $\mathrm{MF}^{gr}_{R,\omega}$ be a **category of a $\mathbb{Z}$-graded matrix factorization** whose object is a matrix factorization $\overline{M} = (M_0, M_1, d_{M_0}, d_{M_1})$ with the potential ω and whose morphism $\overline{f} = (f_0, f_1)$ from $\overline{M} = (M_0, M_1, d_{M_0}, d_{M_1})$ to $\overline{N} = (N_0, N_1, d_{N_0}, d_{N_1})$ consists of a pair of $\mathbb{Z}$-grading preserving R-module morphisms $f_0 : M_0 \to N_0$ and $f_1 : M_1 \to N_1$ such that $d_{N_0} f_0 = f_1 d_{M_0}$ and $d_{N_1} f_1 = f_0 d_{M_1}$. A matrix factorization $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ is **finite** if M_0 and M_1 are finite rank R-modules.

Let R and R' be $\mathbb{Z}$-graded polynomial rings over $\mathbb{Q}$ and S be the maximal ring with inclusions from the ring S to R and R'. We will take the **tensor product** of R and R' over the maximal ring S always,

$$R \underset{S}{\otimes} R' = R \underset{\mathbb{Q}}{\otimes} R' / \{ rs \underset{\mathbb{Q}}{\otimes} r' - r \underset{\mathbb{Q}}{\otimes} sr' | r \in R, r' \in R', s \in S \}.$$

A $\mathbb{Z}$-grading of the tensor product $R \underset{S}{\otimes} R'$ is naturally induced by one of the tensor product $R \underset{\mathbb{Q}}{\otimes} R'$.

For an R-module M and an R'-module M', these tensor product over S is also defined by

$$M \underset{S}{\otimes} N = M \underset{\mathbb{Q}}{\otimes} N / \{ ms \underset{\mathbb{Q}}{\otimes} n - m \underset{\mathbb{Q}}{\otimes} sn | m \in M, n \in N, s \in S \}.$$

Indeed, a $\mathbb{Z}$-grading of the tensor product $M \underset{S}{\otimes} N$ is also induced by one of $M \underset{\mathbb{Q}}{\otimes} N$.

For $\overline{M} = (M_0, M_1, d_{M_0}, d_{M_1}) \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ and $\overline{N} = (N_0, N_1, d_{N_0}, d_{N_1}) \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R',\omega'})$, we define the **tensor product** of matrix factorizations $\overline{M} \boxtimes \overline{N} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R \underset{S}{\otimes} R', \omega + \omega'})$ by

$$\begin{aligned} \overline{M} \boxtimes \overline{N} \quad &:= \quad \left(\begin{pmatrix} M_0 \underset{S}{\otimes} N_0 \\ M_1 \underset{S}{\otimes} N_1 \end{pmatrix}, \begin{pmatrix} M_1 \underset{S}{\otimes} N_0 \\ M_0 \underset{S}{\otimes} N_1 \end{pmatrix}, \begin{pmatrix} d_{M_0} & -d_{N_1} \\ d_{N_0} & d_{M_1} \end{pmatrix}, \begin{pmatrix} d_{M_1} & d_{N_1} \\ -d_{N_0} & d_{M_0} \end{pmatrix} \right), \\ &= \quad \left(\begin{pmatrix} M_0 \underset{S}{\otimes} N_0 \\ M_1 \underset{S}{\otimes} N_1 \end{pmatrix} \xrightarrow[\begin{pmatrix} d_{M_0} & -d_{N_1} \\ d_{N_0} & d_{M_1} \end{pmatrix}]{\qquad\qquad} \begin{pmatrix} M_1 \underset{S}{\otimes} N_0 \\ M_0 \underset{S}{\otimes} N_1 \end{pmatrix} \xrightarrow[\begin{pmatrix} d_{M_1} & d_{N_1} \\ -d_{N_0} & d_{M_0} \end{pmatrix}]{\qquad\qquad} \begin{pmatrix} M_0 \underset{S}{\otimes} N_0 \\ M_1 \underset{S}{\otimes} N_1 \end{pmatrix} \right) \end{aligned}$$

where

$$\begin{pmatrix} d_{M_0} & -d_{N_1} \\ d_{N_0} & d_{M_1} \end{pmatrix} \text{ and } \begin{pmatrix} d_{M_1} & d_{N_1} \\ -d_{N_0} & d_{M_0} \end{pmatrix}$$

simply denote

$$\begin{pmatrix} d_{M_0} \underset{S}{\otimes} \mathrm{Id}_{N_0} & -\mathrm{Id}_{M_1} \underset{S}{\otimes} d_{N_1} \\ \mathrm{Id}_{M_0} \underset{S}{\otimes} d_{N_0} & d_{M_1} \underset{S}{\otimes} \mathrm{Id}_{N_1} \end{pmatrix} \text{ and } \begin{pmatrix} d_{M_1} \underset{S}{\otimes} \mathrm{Id}_{N_0} & \mathrm{Id}_{M_0} \underset{S}{\otimes} d_{N_1} \\ -\mathrm{Id}_{M_1} \underset{S}{\otimes} d_{N_0} & d_{M_0} \underset{S}{\otimes} \mathrm{Id}_{N_1} \end{pmatrix}.$$

This tensor product $\boxtimes$ is commutative and associative. Moreover, there is the unit object for the tensor product.

Proposition 2.1. (1)*For* $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ *and* $\overline{N} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R',\omega'})$, *there is an isomorphism in* $\mathrm{MF}^{gr}_{R \underset{S}{\otimes} R', \omega + \omega'}$

$$\overline{M} \boxtimes \overline{N} \simeq \overline{N} \boxtimes \overline{M}.$$

(2)*For* $\overline{L} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$, $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R',\omega'})$ *and* $\overline{N} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R'',\omega''})$, *there is an isomorphism in* $\mathrm{MF}^{gr}_{R \otimes_S R' \otimes_{S'} R'', \omega+\omega'+\omega''}$

$$(\overline{L} \boxtimes \overline{M}) \boxtimes \overline{N} \simeq \overline{L} \boxtimes (\overline{M} \boxtimes \overline{N}).$$

Proof. See [10]. ☐

Remark 2.2. *As from here,* $\overline{M_1} \boxtimes \overline{M_2} \boxtimes \overline{M_3} \boxtimes \cdots \boxtimes \overline{M_n}$ *is expressed as*

$$\overline{M_1} \boxtimes \overline{M_2} \boxtimes \overline{M_3} \boxtimes \cdots \boxtimes \overline{M_n} = (\cdots((\overline{M_1} \boxtimes \overline{M_2}) \boxtimes \overline{M_3}) \boxtimes \cdots) \boxtimes \overline{M_n}.$$

Proposition 2.3. *The matrix factorization* $(R \xrightarrow{0} 0 \xrightarrow{0} R)$ *is the unit object for tensor product to any object in* $\mathrm{MF}^{gr}_{R,\omega}$. *In brief, for any matrix factorization* $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ *we have*

$$\overline{M} \boxtimes (R \xrightarrow{0} 0 \xrightarrow{0} R) \simeq \overline{M}.$$

Proof. See [10]. ☐

$\overline{R}$ often simply denotes

$$\overline{R} := \ (R \xrightarrow{0} 0 \xrightarrow{0} R)\ .$$

The **translation functor** $\langle 1 \rangle$ changes a matrix factorization $\overline{M} = (M_0, M_1, d_{M_0}, d_{M_1}) \in \mathrm{MF}^{gr}_{R,\omega}$ into

$$\begin{aligned} \overline{M}\,\langle 1 \rangle &= (M_1, M_0, -d_{M_1}, -d_{M_0}) \in \mathrm{MF}^{gr}_{R,\omega} \\ \Big(&= \left(M_1 \xrightarrow{-d_{M_1}} M_0 \xrightarrow{-d_{M_0}} M_1 \right) \Big). \end{aligned}$$

The functor $\langle 2 \rangle$ $(= \langle 1 \rangle^2)$ is the identity functor.

Proposition 2.4. *For* $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ *and* $\overline{N} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R',\omega'})$, *there is an isomorphism in* $\mathrm{MF}^{gr}_{R \otimes_S R', \omega+\omega'}$

$$\begin{aligned} (\overline{M} \boxtimes \overline{N})\,\langle 1 \rangle &= (\overline{M}\,\langle 1 \rangle) \boxtimes \overline{N} \\ &\simeq \overline{M} \boxtimes (\overline{N}\,\langle 1 \rangle). \end{aligned}$$

Proof. See [10]. ☐

The morphism $\overline{f} = (f_1, f_2) : \overline{M} \to \overline{N} \in \mathrm{Mor}(\mathrm{MF}^{gr}_{R,\omega})$ is **null-homotopic** if morphisms $h_0 : M_0 \to N_1$ and $h_1 : M_1 \to N_0$ exist such that $f_0 = h_1 d_{M_0} + d_{N_1} h_0$ and $f_1 = h_0 d_{M_1} + d_{N_0} h_1$. Two morphisms $\overline{f}, \overline{g} : \overline{M} \to \overline{N} \in \mathrm{Mor}(\mathrm{MF}^{gr}_{R,\omega})$ are **homotopic** if $\overline{f} - \overline{g}$ is null-homotopic. In this case, we denote $\overline{f} \sim \overline{g}$. Two matrix factorization $\overline{M}$ and $\overline{N} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ are **homotopy equivalence** if there are $\mathbb{Z}$-grading preserving morphisms $f : \overline{M} \to \overline{N}$ and $g : \overline{N} \to \overline{M}$ such that $fg \sim \mathrm{Id}_{\overline{N}}$ and $gf \sim \mathrm{Id}_{\overline{M}}$.

Let $\mathrm{HMF}^{gr}_{R,\omega}$ be the quotient category of $\mathrm{MF}^{gr}_{R,\omega}$ which has the same objects to $\mathrm{MF}^{gr}_{R,\omega}$ and has morphisms of $\mathrm{Mor}(\mathrm{MF}^{gr}_{R,\omega})$ modulo null-homotopic. The category $\mathrm{HMF}^{gr}_{R,\omega}$ is called the **homotopy category** of $\mathrm{MF}^{gr}_{R,\omega}$. It is obvious that a homotopy equivalence in $\mathrm{MF}^{gr}_{R,\omega}$ is an isomorphism in $\mathrm{HMF}^{gr}_{R,\omega}$.

A matrix factorization in $\mathrm{MF}^{gr}_{R,\omega}$ is called **contractible** if it is isomorphic in $\mathrm{HMF}^{gr}_{R,\omega}$ to the zero matrix factorization

$$\left(0 \longrightarrow 0 \longrightarrow 0 \right).$$

A **$\mathbb{Z}$-grading shift** $\{n\}$ $(n \in \mathbb{Z})$ is an operator up n-grading. That is, for a $\mathbb{Z}$-graded R-module M with the $\mathbb{Z}$-graded decomposition $\oplus M^i$ (M^i:$\mathbb{Q}$ vector space with i-grading), $M\{n\}$ is defined by

$$M\{n\} = \oplus M^{i+n}.$$

The $\mathbb{Z}$-grading shift $\{n\}$ turns the matrix factorization $\overline{M} = (M_0, M_1, d_{M_0}, d_{M_1})$ into

$$\begin{aligned} \overline{M}\{n\} &= (M_0\{n\}, M_1\{n\}, d_{M_0}, d_{M_1}) \\ &= \Big(M_0\{n\} \xrightarrow{d_{M_0}} M_1\{n\} \xrightarrow{d_{M_1}} M_0\{n\} \Big). \end{aligned}$$

Proposition 2.5. *For* $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega})$ *and* $\overline{N} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R',\omega'})$*, there is an equality in* $\mathrm{MF}^{gr}_{R\underset{S}{\otimes}R',\omega+\omega'}$

$$\begin{aligned}(\overline{M}\boxtimes\overline{N})\{n\} &= (\overline{M}\{n\})\boxtimes\overline{N}\\ &= \overline{M}\boxtimes(\overline{N}\{n\})\end{aligned}$$

Proof. We find that these objects are really identical by definition. □

If the potential ω equals 0 then a matrix factorization $\overline{M} \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,0})$ satisfies the boundary condition $d_{M_0}d_{M_1} = d_{M_1}d_{M_0} = 0$. Therefore, there is homology functor H from $\mathrm{MF}^{gr}_{R,0}$ to a category of a $\mathbb{Z}\oplus\mathbb{Z}_2$-graded homology group as follows,

$$\mathrm{H}(\overline{M}) = \bigoplus_{j\in\mathbb{Z},k\in\mathbb{Z}_2}\mathrm{H}^{j,k}(\overline{M}),$$

where k is a complex grading of the matrix factorization $\overline{M}$ (i.e. $k=0$ or 1) and j is a $\mathbb{Z}$-grading derived from the $\mathbb{Z}$-graded modules of the matrix factorization $\overline{M}$. The **Euler characteristic** $\overline{\chi}$ of $H(\overline{M})$ is defined by

$$\overline{\chi}(\mathrm{H}(\overline{M})) = \sum_{j\in\mathbb{Z},k\in\mathbb{Z}_2}\dim_{\mathbb{Q}}\mathrm{H}^{j,k}(\overline{M})q^j.$$

2.2. **Koszul matrix factorization.** Let R be a $\mathbb{Z}$-graded polynomial ring over $\mathbb{Q}$. For homogeneous $\mathbb{Z}$-grading polynomials a, $b\in R$ and a $\mathbb{Z}$-graded R-module M, we define a matrix factorization $K(a;b)_M$ with the potential ab by

$$\begin{aligned}K(a;b)_M &= (M, M\{\frac{1}{2}(\deg(b)-\deg(a))\}, a, b)\\ &= \left(M \xrightarrow{\ a\ } M\{\tfrac{1}{2}(\deg(b)-\deg(a))\} \xrightarrow{\ b\ } M\right),\end{aligned}$$

where $\deg(a)$ and $\deg(b)$ are $\mathbb{Z}$-gradings of the polynomials $a, b\in R$. In general, for sequences $\mathbf{a} = {}^t(a_1,a_2,\ldots,a_k)$, $\mathbf{b} = {}^t(b_1,b_2,\ldots,b_k)$ of homogeneous polynomials in R and R-module M, a matrix factorization $K(\mathbf{a};\mathbf{b})_M$ with the potential $\sum_{i=1}^k a_ib_i$ is defined by

$$K(\mathbf{a};\mathbf{b})_M = \mathop{\boxtimes}_{i=1}^{k} K(a_i;b_i)_R\boxtimes(M,0,0,0).$$

This matrix factorization is called a **Koszul matrix factorization** [5].

Remark 2.6. *Let R be a $\mathbb{Z}$-graded polynomial ring over $\mathbb{Q}$ and let R_y be the $\mathbb{Z}$-graded polynomial ring $R[y]$. For polynomials a and b in R, $K(a;b)_{R_y}$ is a matrix factorization of R_y-modules with rank 1 in $\mathrm{MF}^{gr}_{R_y,ab}$ and a matrix factorization of infinite rank R-modules in $\mathrm{MF}^{gr}_{R,ab}$ besides.*

Proposition 2.7. *Let c be a non-zero element in $\mathbb{Q}$. There is the following isomorphism in $\mathrm{MF}^{gr}_{R,ab}$*

$$K(a;b)_M \simeq K(ca;c^{-1}b)_M.$$

Proof. See [10]. □

Proposition 2.8.

$$\begin{aligned}K(a;b)_M\ \langle 1\rangle &= K(-b;-a)_M\,\{\frac{1}{2}(\deg(b)-\deg(a))\}\\ &\simeq K(b;a)_M\,\{\frac{1}{2}(\deg(b)-\deg(a))\}\end{aligned}$$

Proof. See [10]. □

Proposition 2.9. *Let a_i and b_i be homogeneous $\mathbb{Z}$-grading polynomials such that* $\deg(a_1)+\deg(b_1) = \deg(a_2)+\deg(b_2)$ *and let λ_i $(i=1,2)$ be homogeneous $\mathbb{Z}$-grading polynomials such that* $\deg(\lambda_1) = \deg(a_2)-\deg(a_1)$ *and* $\deg(\lambda_2) = -\deg(b_1)+\deg(a_2)$.
(1) *There is the following isomorphism in* $\mathrm{MF}^{gr}_{R,a_1b_1+a_2b_2}$

$$K\left(\begin{pmatrix}a_1\\a_2\end{pmatrix};\begin{pmatrix}b_1\\b_2\end{pmatrix}\right)_M \simeq K\left(\begin{pmatrix}a_1\\a_2+\lambda_1a_1\end{pmatrix};\begin{pmatrix}b_1-\lambda_1b_2\\b_2\end{pmatrix}\right)_M.$$

(2) There is the following isomorphism in $\mathrm{MF}^{gr}_{R,a_1b_1+a_2b_2}$

$$K\left(\begin{pmatrix} a_1 \\ a_2 \end{pmatrix};\begin{pmatrix} b_1 \\ b_2 \end{pmatrix}\right)_M \simeq K\left(\begin{pmatrix} a_1-\lambda_2 b_2 \\ a_2+\lambda_2 b_1 \end{pmatrix};\begin{pmatrix} b_1 \\ b_2 \end{pmatrix}\right)_M .$$

Proof. See [9][12]. □

Theorem 2.10. *Let* R *be a* $\mathbb{Z}$*-graded polynomial ring and let* a_i, b_i *and* b_i' $(i=1,\cdots,m)$ *be polynomials in* R. *If* $a_1,\cdots,a_m\in R$ *form a regular sequence and*

$$\sum_{i=1}^{m} a_i b_i = \sum_{i=1}^{m} a_i b_i' (=:\omega),$$

there is the following isomorphism in $\mathrm{MF}^{gr}_{R,\omega}$

$$k\left(\begin{pmatrix} a_1 \\ \vdots \\ a_m \end{pmatrix};\begin{pmatrix} b_1 \\ \vdots \\ b_m \end{pmatrix}\right)_M \simeq k\left(\begin{pmatrix} a_1 \\ \vdots \\ a_m \end{pmatrix};\begin{pmatrix} b_1' \\ \vdots \\ b_m' \end{pmatrix}\right)_M .$$

Proof. See Theorem 2.1 on [7] □

Corollary 2.11. *Put* $R=\mathbb{Q}[x_1,x_2,\cdots,x_k]$ *and* $R_y=R[y]\,/\langle y^l+\alpha_1 y^{l-1}+\alpha_2 y^{l-2}+\cdots+\alpha_l\rangle$ $(\alpha_i\in R)$.
(1)Let a_i *be a polynomial* $\in R_y\,(i=1,\cdots,m)$, b_i *be a polynomial* $\in R$ $(i=2,\cdots,m)$ *and let* b_1, β *be polynomials* $\in R_y$ *with the property* $(y+\beta)b_1\in R$. *If these polynomials hold the following conditions;*

(i) $(y+\beta)b_1$, b_2, $\cdots$, b_m *form a regular sequence in* R,

(ii) $a_1b_1(y+\beta)+\sum_{i=2}^{m} a_i b_i$ $(=:\omega)\in R$,

then there are polynomials $a_i'\in R$ $(i=1,\cdots,m)$ *to give the following isomorphism in* $\mathrm{MF}_{R,\omega}$

$$K\left(\begin{pmatrix} (y+\beta)a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix};\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} \simeq K\left(\begin{pmatrix} (y+\beta)a_1' \\ a_2' \\ \vdots \\ a_m' \end{pmatrix};\begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

(2)Let a_i *be a polynomial* $\in R_y$ $(i=1,\cdots,m)$, b_i *be a polynomial* $\in R$ $(i=1,\cdots,m)$ *and* β *be a polynomial* $\in R$. *If these polynomials hold the following conditions;*

(i) b_1, b_2, $\cdots$, b_m *form a regular sequence in* R,

(ii) $a_1b_1(y+\beta)+\sum_{i=2}^{m} a_i b_i$ $(=:\omega')\in R$,

then there are polynomials $a_1'\in R_y$ *and* $a_i'\in R$ $(i=2,\cdots,m)$ *to give the following isomorphism in* $\mathrm{MF}_{R,\omega'}$

$$K\left(\begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix};\begin{pmatrix} b_1(y+\beta) \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} \simeq K\left(\begin{pmatrix} a_1' \\ a_2' \\ \vdots \\ a_m' \end{pmatrix};\begin{pmatrix} b_1(y+\beta) \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

Proof. (1)We consider the matrix factorization

$$K\left(\begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix};\begin{pmatrix} (y+\beta)b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

Since the sequence $((y+\beta)b_1, b_2, \cdots, b_m)$ is regular, by Corollary 2.10 there are polynomials $a_i' \in R$ $(i = 1, \cdots, m)$ to give the following matrix factorization,

$$K\left(\begin{pmatrix} a_1' \\ a_2' \\ \vdots \\ a_m' \end{pmatrix} ; \begin{pmatrix} (y+\beta)b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

It is obvious that the polynomial degree of b_1 is less than one of $(y+\beta)b_1$. Then we have the isomorphism

$$K\left(\begin{pmatrix} (y+\beta)a_1 \\ a_2 \\ \vdots \\ a_m \end{pmatrix} ; \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} \simeq K\left(\begin{pmatrix} (y+\beta)a_1' \\ a_2' \\ \vdots \\ a_m' \end{pmatrix} ; \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

(2)The polynomial a_1 can be described as

$$a_1 = \gamma_0 y^{l-1} + \gamma_1 y^{l-2} + \cdots + \gamma_{l-1} \qquad (\gamma_i \in R). \tag{1}$$

First we consider the matrix factorization

$$K\left(\begin{pmatrix} a_1(y+\beta) \\ a_2 \\ \vdots \\ a_m \end{pmatrix} ; \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

By assumption, the second sequence $(b_1, b_2, \cdots, b_m)$ is regular and potential of the matrix factorization is in R. Using Theorem 2.10 we find that there are polynomials c_1 and a_i' $(i = 2, \cdots, m)$ in R to give an isomorphism between the above matrix factorization and the following matrix factorization

$$K\left(\begin{pmatrix} c_1 \\ a_2' \\ \vdots \\ a_m' \end{pmatrix} ; \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}\right)_{R_y} .$$

Therefore, $a_1(y+\beta) - c_1$ can be described as an R_y coefficient linear sum of b_2, $\cdots$, b_{m-1}, b_m. Since, by Equation (1),

$$\begin{aligned} a_1(y+\beta) &= \gamma_0 y^l + (\gamma_1 + \beta\gamma_0) y^{l-1} + (\gamma_2 + \beta\gamma_1) y^{l-2} + \cdots + (\gamma_{l-1} + \beta\gamma_{l-2}) y + \beta\gamma_{l-1} \\ &= (\gamma_1 + \beta\gamma_0 - \alpha_1\gamma_0) y^{l-1} + \cdots + (\gamma_{l-1} + \beta\gamma_{l-2} - \alpha_{l-1}\gamma_0) y + \beta\gamma_{l-1} - \alpha_l\gamma_0 \end{aligned}$$

and b_2, $\cdots$, b_{m-1}, b_m form a regular sequence, the polynomials $\gamma_i + \beta\gamma_{i-1} - \alpha_i\gamma_0$ $(i = 1, \cdots, l-1)$ are described as an R coefficient linear sum of b_2, $\cdots$, b_{m-1}, b_m,

$$\gamma_i + \beta\gamma_{i-1} - \alpha_i\gamma_0 = \sum_{j=2}^{m} s_{i,j} b_j \qquad (i = 1, \cdots, l-1), \tag{2}$$

and $\beta\gamma_{l-1} - \alpha_l\gamma_0$ is described as

$$\beta\gamma_{l-1} - \alpha_l\gamma_0 = c_1 + \sum_{j=2}^{m} s_{l,j} b_j .$$

Moreover equations of (2) change into the following equations by linear transform,

$$\gamma_i = \left(\sum_{j=0}^{i} (-1)^{i-j} \alpha_j \beta^{i-j}\right)\gamma_0 + \sum_{j=2}^{m} \widetilde{s_{i,j}} b_j \qquad (i = 1, \cdots, l-1),$$

where $\alpha_0 = 1$ and $\widetilde{s_{i,j}} = \sum_{k=1}^{i} (-1)^k \beta^k s_{i-k,j}$.

Thus we have the equivalence between matrix factorizations as follows,

$$K\left(\left(\begin{array}{c} a_1 \\ a_2 \\ \vdots \\ a_m \end{array}\right); \left(\begin{array}{c} b_1(y+\beta) \\ b_2 \\ \vdots \\ b_m \end{array}\right)\right)_{R_y} \simeq K\left(\left(\begin{array}{c} a_1{}' \\ a_2 - \sum_{i=1}^{l-1} \widetilde{s_{i,2}} b_1 y^{l-1-i} \\ \vdots \\ a_m - \sum_{i=1}^{l-1} \widetilde{s_{i,m}} b_1 y^{l-1-i} \end{array}\right); \left(\begin{array}{c} b_1(y+\beta) \\ b_2 \\ \vdots \\ b_m \end{array}\right)\right)_{R_y},$$

where

$$a_1{}' = \sum_{i=0}^{l-1} \left(\sum_{j=0}^{i} (-1)^{i-j} \alpha_j \beta^{i-j} \right) \gamma_0 y^{l-1-i}.$$

It is obvious to find the polynomial $a_1{}'(y+\beta)$ is in R. Since the polynomials b_2, $\cdots$, b_{m-1}, b_m form a regular sequence and the polynomial $\sum_{j=2}^{m}(a_j - \sum_{i=1}^{l-1} \widetilde{s_{i,j}} b_1 y^{l-1-i}) b_j = \omega' - a_1{}'(y+\beta)$ is in R, polynomials $a_i{}' \in R$ $(i = 2, \cdots, m)$ exist to give the following isomorphism

$$K\left(\left(\begin{array}{c} a_1{}' \\ a_2 - \sum_{i=1}^{l-1} \widetilde{s_{i,2}} b_1 y^{l-1-i} \\ \vdots \\ a_m - \sum_{i=1}^{l-1} \widetilde{s_{i,m}} b_1 y^{l-1-i} \end{array}\right); \left(\begin{array}{c} b_1(y+\beta) \\ b_2 \\ \vdots \\ b_m \end{array}\right)\right)_{R_y} \simeq K\left(\left(\begin{array}{c} a_1{}' \\ a_2{}' \\ \vdots \\ a_m{}' \end{array}\right); \left(\begin{array}{c} b_1(y+\beta) \\ b_2 \\ \vdots \\ b_m \end{array}\right)\right)_{R_y}.$$

□

Remark 2.12. *Put* $R_y = R[y]/\langle y^l + \alpha_1 y^{l-1} + \alpha_2 y^{l-2} + \cdots + \alpha_l \rangle$ $(\alpha_i \in R)$*. If the variable* y *remains in a polynomial* p *then the multiplication map* p *to* R_y *is complicated as an* R*-module morphism. However, if the variable* y *dose not exist in a polynomial* p *then the multiplication map* p *is simply a diagonal map as an* R*-module morphism,*

$$R_y \xrightarrow{p} R_y = \left(\begin{array}{r} \alpha_0 R \\ \alpha_1 R \\ \vdots \\ \alpha_{l-1} R \end{array}\right) \xrightarrow[\left(\begin{array}{cccc} p & 0 & \cdots & 0 \\ 0 & p & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & p \end{array}\right)]{} \left(\begin{array}{r} \alpha_0 R \\ \alpha_1 R \\ \vdots \\ \alpha_{l-1} R \end{array}\right),$$

where α_0, α_1, $\cdots$, α_{l-1} *form a basis of* R_y *as an* R*-module.*

In next section, Corollary 2.11 is useful for decomposing a matrix factorization into its direct sum of matrix factorizations.

The following theorem is a generalization to multivariable of Theorem 2.2 given by Khovanov and Rozansky [7][10].

Theorem 2.13. *We put* $R = \mathbb{Q}[\underline{x}]$ *and* $R_{\mathbf{y}} = R[\underline{y}]$*, where* $\underline{x} = (x_1, x_2, \cdots, x_l)$ *and* $\underline{y} = (y_1, y_2, \cdots, y_m)$*. If* $\mathbf{a} = {}^t(a_1, a_2, \cdots, a_k)$ *and* $\mathbf{b} = {}^t(b_1, b_2, \cdots, b_k)$*, where* a_i *and* b_i *are homogeneous polynomials in* $R_{\mathbf{y}}$*, satisfy the following conditions*

(i) $\sum_{i=1}^{k} a_i b_i \,(=: \omega) \in R$,

(ii) *There exists* b_j *which can be described by* $c y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} + p$*, where* c *is a constant and* p *is a polynomial of* $R_{\mathbf{y}}$ *whose every monomials are not zero in the quotient ring* $R_{\mathbf{y}}/\langle y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} \rangle$*,*

then there is the following isomorphism in $\mathrm{HMF}^{gr}_{R,\omega}$*,*

$$K(\mathbf{a};\mathbf{b})_{R_{\mathbf{y}}} \simeq K(\overset{j}{\check{\mathbf{a}}};\overset{j}{\check{\mathbf{b}}})_{R_{\mathbf{y}}/\langle b_j \rangle},$$

where $\overset{j}{\check{\mathbf{a}}}$ *and* $\overset{j}{\check{\mathbf{b}}}$ *are the sequences omitted the* i*-th polynomial of* $\mathbf{a}$ *and* $\mathbf{b}$.

Proof. We can prove this theorem to repeat proof by M.Khovanov and L.Rozansky in [7][10]. By Proposition 2.7 and the above assumption, we can describe the left-hand matrix factorization

$$K(\mathbf{a};\mathbf{b})_{R_{\mathbf{y}}} \simeq K(\overset{i}{\check{\mathbf{a}}};\overset{i}{\check{\mathbf{b}}})_{R_{\mathbf{y}}} \boxtimes K(a_j; y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} + p)_{R_{\mathbf{y}}}.$$

The Koszul matrix factorization $K(\overset{i}{\check{\mathbf{a}}};\overset{i}{\check{\mathbf{b}}})_{R_{\mathbf{y}}}$ is described as $(R_{\mathbf{y}}^r, R_{\mathbf{y}}^r, D_0, D_1)$, where D_0, $D_1 \in \mathrm{Mat}_r(R_{\mathbf{y}})$ ($r = 2^{k-2}$). Then we have

$$K(\mathbf{a};\mathbf{b})_{R_{\mathbf{y}}} = \left(\begin{array}{ccccc} R_{\mathbf{y}}^r & & R_{\mathbf{y}}^r & & R_{\mathbf{y}}^r \\ \oplus & \xrightarrow{D_0{\acute{}}} & \oplus & \xrightarrow{D_1{\acute{}}} & \oplus \\ R_{\mathbf{y}}^r & & R_{\mathbf{y}}^r & & R_{\mathbf{y}}^r \end{array} \right),$$

where $D_0{\acute{}} = \begin{pmatrix} D_0 & (-y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} - p)\mathrm{Id}_{R_{\mathbf{y}}^r} \\ a_j \mathrm{Id}_{R_{\mathbf{y}}^r} & D_1 \end{pmatrix}$, $D_1{\acute{}} = \begin{pmatrix} D_1 & (y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} + p)\mathrm{Id}_{R_{\mathbf{y}}^r} \\ -a_j \mathrm{Id}_{R_{\mathbf{y}}^r} & D_0 \end{pmatrix}$.

The ring $R_{\mathbf{y}}$ is split into the direct sum as an R-module;

$$R_{\mathbf{y}} \simeq R_< \oplus R_\geq,$$

where

$$R_< = \bigoplus_{\substack{(i_1, i_2, \cdots, i_m) \in \mathbb{N}_{\geq 0}^m \\ \min\{i_1 - n_1, i_2 - n_2, \cdots, i_m - n_m\} < 0}} R \cdot y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} \simeq R_{\mathbf{y}} / \left\langle y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} + p \right\rangle$$

and

$$R_\geq = \bigoplus_{(i_1, i_2, \cdots, i_m) \in \mathbb{N}_{\geq 0}^m} R \cdot y_1^{i_1} y_2^{i_2} \cdots y_m^{i_m} (y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} + p).$$

The R-module morphism $R_{\mathbf{y}} \xrightarrow{y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} + p} R_{\mathbf{y}}$ induces the R-module isomorphism

$$f_{iso} : R_{\mathbf{y}} \xrightarrow{y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} + p} R_\geq.$$

Moreover, there are the natural R-module injection

$$f_{inj} : R_< \xrightarrow{} R_{\mathbf{y}},$$

and the natural R-module projections

$$f_{proj_<} : R_{\mathbf{y}} \xrightarrow{} R_< \ ,$$

$$f_{proj_\geq} : R_{\mathbf{y}} \xrightarrow{} R_\geq.$$

The R-module $R_{\mathbf{y}}^r$ is also split into the direct sum

$$R_{\mathbf{y}}^r \simeq R_<^r \oplus R_\geq^r.$$

Then there are also the R-module isomorphism

$$F_{iso} : R_{\mathbf{y}}^r \xrightarrow{y_1^{n_1} y_2^{n_2} \cdots y_m^{n_m} + p} R_\geq^r,$$

the R-module injection

$$F_{inj} : R_<^r \xrightarrow{} R_{\mathbf{y}}^r$$

and the R-module projections

$$F_{proj_<} : R_{\mathbf{y}}^r \xrightarrow{} R_<^r \ ,$$

$$F_{proj_\geq} : R_{\mathbf{y}}^r \xrightarrow{} R_\geq^r.$$

Since the following R-module morphisms ϕ_0 and ϕ_1 are R-module isomorphisms;

$$\phi_0 = \begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \xrightarrow[\begin{pmatrix} F_{inj} & (y_1^{n_1}y_2^{n_2}\cdots y_m^{n_m}+p)F_{iso}^{-1} & 0 \\ 0 & D_0\,F_{iso}^{-1} & \mathrm{Id}_{R^r_{\mathbf{y}}} \end{pmatrix}]{\hspace{8cm}} \begin{matrix} R^r_{\mathbf{y}} \\ \oplus \\ R^r_{\mathbf{y}}, \end{matrix}$$

$$\phi_1 = \begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \xrightarrow[\begin{pmatrix} F_{inj} & (-y_1^{n_1}y_2^{n_2}\cdots y_m^{n_m}-p)F_{iso}^{-1} & 0 \\ 0 & D_1\,F_{iso}^{-1} & \mathrm{Id}_{R^r_{\mathbf{y}}} \end{pmatrix}]{\hspace{8cm}} \begin{matrix} R^r_{\mathbf{y}} \\ \oplus \\ R^r_{\mathbf{y}}, \end{matrix}$$

the Koszul matrix factorization $K(\mathbf{a};\mathbf{b})_{R_{\mathbf{y}}}$ is isomorphic to the following matrix factorization $\overline{M}$ by the isomorphism $\overline{\phi}=(\phi_0,\phi_1)$,

$$\overline{M} = \left(\begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \xrightarrow[\phi_1^{-1}\,D_0'\phi_0]{\hspace{3cm}} \begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \xrightarrow[\phi_0^{-1}\,D_1'\phi_1]{\hspace{3cm}} \begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \right).$$

Since $\overline{\phi}^{-1}$ consists of

$$\phi_0^{-1} = \begin{pmatrix} F_{proj_<} & 0 \\ (y_1^{n_1}y_2^{n_2}\cdots y_m^{n_m}+p)F_{iso}^{-1}\,F_{proj_\geq} & 0 \\ -D_0F_{iso}^{-1}\,F_{proj_\geq} & \mathrm{Id}^r_{R_{\mathbf{y}}} \end{pmatrix} \text{ and } \phi_1^{-1} = \begin{pmatrix} F_{proj_<} & 0 \\ (-y_1^{n_1}y_2^{n_2}\cdots y_m^{n_m}-p)F_{iso}^{-1}\,F_{proj_\geq} & 0 \\ D_1F_{iso}^{-1}\,F_{proj_\geq} & \mathrm{Id}^r_{R_{\mathbf{y}}} \end{pmatrix},$$

the morphisms $\phi_1^{-1}\,D_0'\phi_0$ and $\phi_0^{-1}\,D_1'\phi_1$ are described by

$$\phi_1^{-1}\,D_0'\phi_0 = \begin{pmatrix} F_{proj_<}\,D_0\,F_{inj} & 0 & 0 \\ G_1 & 0 & F_{iso} \\ G_2 & \omega F_{iso}^{-1} & 0 \end{pmatrix} \text{ and } \phi_0^{-1}\,D_1'\phi_1 = \begin{pmatrix} F_{proj_<}\,D_1\,F_{inj} & 0 & 0 \\ H_1 & 0 & F_{iso} \\ H_2 & \omega F_{iso}^{-1} & 0 \end{pmatrix}.$$

By the construction we have

$$\begin{aligned} G_1\,F_{proj_<}\,D_1\,F_{inj} + F_{iso}\,H_2 &= 0, \\ G_2\,F_{proj_<}\,D_1\,F_{inj} + \omega F_{iso}^{-1}\,H_1 &= 0, \\ H_1\phi_1^{-1}\,D_0'\phi_0 + F_{iso}G_2 &= 0, \\ H_2\phi_1^{-1}\,D_0'\phi_0 + \omega F_{iso}^{-1}G_1 &= 0. \end{aligned}$$

By the isomorphism $\overline{\psi} = \left(\begin{pmatrix} \mathrm{Id}_{R^r_<} & 0 & 0 \\ 0 & \mathrm{Id}_{R^r_\geq} & 0 \\ F_{iso}^{-1}G_1 & 0 & \mathrm{Id}_{R^r_{\mathbf{y}}} \end{pmatrix}, \begin{pmatrix} \mathrm{Id}_{R^r_<} & 0 & 0 \\ 0 & \mathrm{Id}_{R^r_\geq} & 0 \\ F_{iso}^{-1}H_1 & 0 & \mathrm{Id}_{R^r_{\mathbf{y}}} \end{pmatrix} \right)$, the matrix factorization $\overline{M}$ is isomorphic to the following matrix factorization,

$$\overline{M} = \left(\begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \xrightarrow[\begin{pmatrix} \phi_1^{-1}\,D_0'\phi_0 & 0 & 0 \\ 0 & 0 & F_{iso} \\ 0 & \omega F_{iso}^{-1} & 0 \end{pmatrix}]{\hspace{5cm}} \begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \xrightarrow[\begin{pmatrix} \phi_1^{-1}\,D_1'\phi_0 & 0 & 0 \\ 0 & 0 & F_{iso} \\ 0 & \omega F_{iso}^{-1} & 0 \end{pmatrix}]{\hspace{5cm}} \begin{matrix} R^r_< \\ \oplus \\ R^r_\geq \\ \oplus \\ R^r_{\mathbf{y}} \end{matrix} \right).$$

The submatrix factorization

$$\left(\begin{pmatrix} R^r_{\geq}\\ \oplus \\ R^r_{\mathbf{y}}\end{pmatrix} \xrightarrow[\begin{pmatrix}0 & F_{iso}\\ \omega F_{iso}^{-1} & 0\end{pmatrix}]{} \begin{pmatrix} R^r_{\geq}\\ \oplus \\ R^r_{\mathbf{y}}\end{pmatrix} \xrightarrow[\begin{pmatrix}0 & F_{iso}\\ \omega F_{iso}^{-1} & 0\end{pmatrix}]{} \begin{pmatrix} R^r_{\geq}\\ \oplus \\ R^r_{\mathbf{y}}\end{pmatrix}\right)$$

is contractible in $\mathrm{HMF}^{gr}_{R,\omega}$. Thus, $\overline{M}$ is isomorphic to the following matrix factorization

$$\left(R^r_{<} \xrightarrow{F_{proj_<}\, D_0\, F_{inj}} R^r_{<} \xrightarrow{F_{proj_<}\, D_1\, F_{inj}} R^r_{<}\right)$$

in $\mathrm{HMF}^{gr}_{R,\omega}$. By the choice of a basis of $R_{\mathbf{y}}$ as an R-module, we find that this matrix factorization is isomorphic to $K(\overset{j}{\check{\mathbf{a}}};\overset{j}{\check{\mathbf{b}}})_{R_{\mathbf{y}}/\langle b_j\rangle}$. ☐

Corollary 2.14. *We put $R=\mathbb{Q}[\underline{x}]$ and $R_{\mathbf{y}}=R[\underline{y}]$, where $\underline{x}=(x_1,x_2,\cdots,x_l)$ and $\underline{y}=(y_1,y_2,\cdots,y_m)$. If $\mathbf{a}={}^t(a_1,a_2,\cdots,a_k)$ and $\mathbf{b}={}^t(b_1,b_2,\cdots,b_k)$, where $a_i,b_i\in R_{\mathbf{y}}$, satisfy the following conditions,*

(i) $\sum_{i=1}^{k} a_ib_i\ (=:\omega)\in R$,

($*$) *There exist a polynomial $b_j\in R_{\mathbf{y}}$ such that ,when $b_j(\underline{x},\underline{y})$ denotes $b_j(\underline{0},\underline{y})+p$ ($p\in R_{\mathbf{y}}$), every monomial of p is not zero in the quotient ring $R_{\mathbf{y}}/\left\langle b_j(\underline{0},\underline{y})\right\rangle$,*

then there is the following isomorphism in $\mathrm{HMF}^{gr}_{R,\omega}$,

$$K(\mathbf{a};\mathbf{b})_{R_{\mathbf{y}}}\simeq K(\overset{j}{\check{\mathbf{a}}};\overset{j}{\check{\mathbf{b}}})_{R_{\mathbf{y}}/\langle b_j\rangle}.$$

Proof. Each monomial of $b_j(\underline{0},\underline{y})$ is suitable for $y_1^{i_1}y_2^{i_2}\cdots y_m^{i_m}$ of the condition (ii) of Theorem 2.13. Thus, we obtain this corollary. ☐

Corollary 2.15. *We put $R=\mathbb{Q}[\underline{x}]$ and $R_{\mathbf{y}}=R[\underline{y}]$, where $\underline{x}=(x_1,x_2,\cdots,x_l)$ and $\underline{y}=(y_1,y_2,\cdots,y_m)$. If $\mathbf{a}={}^t(a_1,a_2,\cdots,a_k)$ and $\mathbf{b}={}^t(b_1,b_2,\cdots,b_k)$, where $a_i,b_i\in R_{\mathbf{y}}$, satisfy the following conditions,*

(i) $\sum_{i=1}^{k} a_ib_i\ (=:\omega)\in R$,

(ii) *There are polynomials $b_{j_1},b_{j_2},\cdots,b_{j_l}$ holding the condition ($*$) of Corollary 2.14 and the condition that the sequence*

$$(b_{j_1}(\underline{0},\underline{y}),b_{j_2}(\underline{0},\underline{y}),\cdots,b_{j_l}(\underline{0},\underline{y}))$$

forms regular,

then there is the following isomorphism in $\mathrm{HMF}^{gr}_{R,\omega}$,

$$K(\mathbf{a};\mathbf{b})_{R_{\mathbf{y}}}\simeq K(\overset{j_1,j_2,\cdots,j_l}{\check{\mathbf{a}}};\overset{j_1,j_2,\cdots,j_l}{\check{\mathbf{b}}})_{R_{\mathbf{y}}/\langle b_{j_1},b_{j_2},\cdots,b_{j_l}\rangle}.$$

Proof. Since the sequence $(b_{j_1}(\underline{0},\underline{y}),b_{j_2}(\underline{0},\underline{y}),\cdots,b_{j_l}(\underline{0},\underline{y}))$ is regular. Then the sequences $\overset{j_l}{\check{\mathbf{a}}}$ and $\overset{j_l}{\check{\mathbf{b}}}$ satisfy the conditions (i) and (ii) after we apply Corollary 2.14 to the polynomial $b_{j_l}(\underline{0},\underline{y})$. Thus we can prove this corollary using induction. ☐

3. Matrix factorizations for colored planar diagrams

3.1. **Poincaré series of a $\mathbb{Z}$-graded algebra.** We briefly recall the Poincaré series of a $\mathbb{Z}$-graded algebra. We only state its definition and properties.

Let $R=\bigoplus_{i=0}^{\infty}R_i$ be a $\mathbb{Z}$-graded algebra over $\mathbb{Q}$. The **Poincaré series** of R is defined to be

$$P_q(R):=\sum_{i=0}^{\infty}(\dim_{\mathbb{Q}}R_i)q^i.$$

If x is a homogeneous element with n-grading in R and not a zero divisor, then we find

$$P_q(R/xR) = P_q(R)(1-q^n).$$

Generally, for a regular sequence $\mathcal{X}_r = (x_1, x_2, \cdots, x_r)$ whose each variable x_i has homogeneous n_i-gradings respectively, we find

$$P_q(R/\langle \mathcal{X}_r\rangle) = P_q(R)(1-q^{n_1})(1-q^{n_2})\cdots(1-q^{n_r}), \tag{3}$$

where $\langle \mathcal{X}_r\rangle$ is an ideal generated by $x_1, x_2, \cdots, x_r$.

Proposition 3.1. *The homogeneous* $\mathbb{Z}$*-grading terms of*

$$(1 + x_1 + x_2 + \cdots + x_r)(1 + y_1 + y_2 + \cdots + y_s) - 1,$$

where $\deg x_i = 2i$ *and* $\deg y_i = 2i$*, form a regular sequence in* $\mathbb{Q}[x_1, x_2, \cdots, x_r, y_1, y_2, \cdots, y_s]$.

Let I be an ideal generated by homogeneous $\mathbb{Z}$-grading terms of

$$(1 + x_1 + x_2 + \cdots + x_r)(1 + y_1 + y_2 + \cdots + y_s) - 1,$$

Since these homogeneous terms form a regular sequence, by this proposition and the equation (3) the Poincaré series of R/I is

$$P_q(R/I) = \frac{(1-q^2)(1-q^4)\cdots(1-q^{2r+2s})}{(1-q^2)(1-q^4)\cdots(1-q^{2r})(1-q^2)(1-q^4)\cdots(1-q^{2s})}. \tag{4}$$

Let x_i be the elementary symmetric functions in r variables $t_1, t_2, \cdots, t_r$ (a $\mathbb{Z}$-grading of every variable t_i is 2);

$$x_i = \sum_{1\leq n_1<n_2<\cdots<n_i\leq r} t_{n_1} t_{n_2} \cdots t_{n_i} \tag{5}$$

and let F_r be an i variables function obtained by expanding the power sum $t_1^{n+1} + t_2^{n+1} + \cdots + t_r^{n+1}$ with the elementary symmetric functions $x_1, x_2, \cdots, x_r$, i.e.

$$F_i(x_1, x_2, \cdots, x_i) = t_1^{n+1} + t_2^{n+1} + \cdots + t_i^{n+1}. \tag{6}$$

The $\mathbb{Z}$-grading of x_i is $2i$ and one of F_r is $2n+2$. Using a sequence of i variables $\mathcal{X}_i = (x_1, x_2, \cdots, x_i)$ We often denotes $F_i(x_1, x_2, \cdots, x_i)$ by $F_i(\mathcal{X}_i)$.

We define **Jacobi algebra** J_{F_r} to be

$$J_{F_r} = \mathbb{Q}[x_1, x_2, \cdots, x_r]/\left\langle \frac{\partial F_r}{\partial x_1}, \frac{\partial F_r}{\partial x_2}, \cdots, \frac{\partial F_r}{\partial x_r}\right\rangle.$$

In the work of Gepner [2], the Jacobi algebra J_{F_r} is isomorphic to the cohomology ring of the Grassmannian manifold $\mathrm{H}^*(Gr_r(\mathbb{C}^n))$ as a $\mathbb{Z}$-graded algebra. Since the Poincaré series of the cohomology ring $\mathrm{H}^*(Gr_r(\mathbb{C}^n))$ is

$$\frac{(1-q^2)(1-q^4)\cdots(1-q^{2n})}{(1-q^2)(1-q^4)\cdots(1-q^{2r})(1-q^2)(1-q^4)\cdots(1-q^{2n-2r})},$$

the Poincaré series of Jacobi algebra J_{F_r} is also

$$P_q(J_{F_r}) = \frac{(1-q^2)(1-q^4)\cdots(1-q^{2n})}{(1-q^2)(1-q^4)\cdots(1-q^{2r})(1-q^2)(1-q^4)\cdots(1-q^{2n-2r})}. \tag{7}$$

3.2. **Colored planar diagrams and matrix factorizations.** We define matrix factorizations for the colored planar diagrams in Figure 5. Let $t_{i,k}$ be a variable with $\mathbb{Z}$-grading 2 for any formal indexes i and k, let $x_{j,k}$ be the elementary symmetric function with $\mathbb{Z}$-grading $2j$ in a polynomial ring $\mathbb{Q}[t_{1,k}, t_{2,k}, \cdots, t_{i,k}]$, where k is a formal index, and let F_i be an i variables function obtained by expanding the power sum $t_{1,k}^{n+1} + t_{2,k}^{n+1} + \cdots + t_{i,k}^{n+1}$ with the elementary symmetric function $x_{1,k}, x_{2,k}, \cdots, x_{i,k}$ (as Equation (6)).

We assign a sequence $\mathcal{X}_{i,k}$ of i variables $x_{1,k}$, $x_{2,k}$, $\cdots$, $x_{i,k}$ on a boundary of a line colored i and define a map $\mathcal{C}$ from a colored planar diagram with such an assignment to a matrix factorization.

Definition 3.2. *A matrix factorization for*



(i = 1, $\cdots$, n)

is defined to be



$$\mathcal{C}\left(\qquad \right)_n := \mathop{\boxtimes}_{j=1}^{i} K\left(L^{1;2}_{j,i}; x_{j,1}-x_{j,2}\right)_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]}, \tag{8}$$

where

$$L^{1;2}_{j,i} = \frac{F_i(x_{1,2},\cdots,x_{j-1,2},x_{j,1},x_{j+1,1},\cdots,x_{i,1}) - F_i(x_{1,2},\cdots,x_{j-1,2},x_{j,2},x_{j+1,1},\cdots,x_{i,1})}{x_{j,1}-x_{j,2}}$$

and

$$\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}] = \mathbb{Q}[x_{1,1},\cdots,x_{i,1},x_{1,2},\cdots,x_{i,2}].$$

This matrix factorization is an object of $\mathrm{MF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}],F_i(\mathcal{X}_{i,1})-F_i(\mathcal{X}_{i,2})}$.

Remark 3.3. *For $i \geq n+1$, we can also consider the matrix factorization for a line colored i,* *as the above definition. However, we find that these matrix factorizations are homotopic to the zero matrix factorization.*

Definition 3.4. *A matrix factorization for the following trivalent diagrams,*



and



,

is defined to be



$$\mathcal{C}\left(\qquad \right)_n := \mathop{\boxtimes}_{j=1}^{i_3} K\left(\Lambda^{1;2,3}_{j,i_1;i_2}; x_{j,3}-X^{1;2}_{j,i_1;i_2}\right)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3}]}, \tag{9}$$



$$\mathcal{C}\left(\qquad \right)_n := \mathop{\boxtimes}_{j=1}^{i_3} K\left(V^{1;2,3}_{j,i_1;i_2}; X^{1;2}_{j,i_1;i_2}-x_{j,3}\right)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3}]}\{-i_1 i_2\}, \tag{10}$$

where

$$\Lambda^{1;2,3}_{j,i_1;i_2} = \frac{F_{i_3}(X^{1;2}_{1,i_1;i_2},\cdots,X^{1;2}_{j-1,i_1;i_2},x_{j,3},x_{j+1,3},\cdots,x_{i_3,3}) - F_{i_3}(X^{1;2}_{1,i_1;i_2},\cdots,X^{1;2}_{j-1,i_1;i_2},X^{1;2}_{j,i_1;i_2},x_{j+1,3},\cdots,x_{i_3,3})}{x_{j,3}-X^{1;2}_{j,i_1;i_2}},$$

$$V^{1;2,3}_{j,i_1;i_2} = \frac{F_{i_3}(x_{1,3},\cdots,x_{j-1,3},X^{1;2}_{j,i_1;i_2},X^{1;2}_{j+1,i_1;i_2},\cdots,X^{1;2}_{i_3,i_1;i_2}) - F_{i_3}(x_{1,3},\cdots,x_{j-1,3},x_{j,3},X^{1;2}_{j+1,i_1;i_2},\cdots,X^{1;2}_{i_3,i_1;i_2})}{X^{1;2}_{j,i_1;i_2}-x_{j,3}},$$

and $X^{k_1;k_2}_{j,i_1;i_2}$ is the $2j$-grading term of $(1+x_{1,k_1}+x_{2,k_1}+\cdots+x_{i_1,k_1})(1+x_{1,k_2}+x_{2,k_2}+\cdots+x_{i_2,k_2})-1$.

Two matrix factorizations (9) and (10) are an object of $\mathrm{MF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3}],F_{i_3}(\mathcal{X}_{i_3,3})-F_{i_1}(\mathcal{X}_{i_1,1})-F_{i_2}(\mathcal{X}_{i_2,2})}$ and an object of $\mathrm{MF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3}],F_{i_1}(\mathcal{X}_{i_1,1})+F_{i_2}(\mathcal{X}_{i_2,2})-F_{i_3}(\mathcal{X}_{i_3,3})}$ respectively.

We define a matrix factorization for a more general planar diagram by using tensor product as follows.

We consider two planar diagrams which have a line colored i and can be match with keeping the orientation on the line colored i,



and



.

Using the matrix factorizations

$\mathcal{C}\Big($  $\Big)_n \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R,\omega+F_i(\mathcal{X}_{i,1})})$ and $\mathcal{C}'\Big($  $\Big)_n \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R',\omega'-F_i(\mathcal{X}_{i,2})})$,

we define a matrix factorization for the glued planar diagram

$\mathcal{C}'\Big($  $\Big)_n \in \mathrm{Ob}(\mathrm{MF}^{gr}_{R\underset{S}{\otimes}R',\omega+\omega'})$ by $\mathcal{C}'\Big($  $\Big)_n \boxtimes \mathcal{C}'\Big($  $\Big)_n\Big|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}$.

This means that we identify the sequence $\mathcal{X}_{i,1}$ and the sequence $\mathcal{X}_{i,2}$ after taking the tensor product of these matrix factorizations. Since a potential of the tensor product of two matrix factorizations is the sum of each potential, the potential of the glued matrix factorization is $\omega+\omega'$.

For a planar diagram Γ composed of the disjoint union of planar diagrams Γ_1 and Γ_2, we define

$$\mathcal{C}'(\ \Gamma\ )_n := \mathcal{C}'(\ \Gamma_1\ )_n \boxtimes \mathcal{C}'(\ \Gamma_2\ )_n$$

Since we only consider a colored planar diagram decomposing into the colored trivalent diagrams in Figure 5, we can obtain a matrix factorization for the colored planar diagram by taking tensor product of matrix factorizations for the trivalent diagrams. By definition of gluing, the potential of a colored planar diagram depend only on the boundary assignment of colored planar diagram.

Proposition 3.5. **(1)** *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i,1}]\underset{S}{\otimes}R,\omega+F_i(\mathcal{X}_{i,1})}$, *where the polynomial ring R and the potential ω are determined by the boundary sequences except the sequence $\mathcal{X}_{i,1}$,*

$\mathcal{C}'\Big($  $\Big)_n \simeq \mathcal{C}'\Big($  $\Big)_n$

(2) *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i,1}]\underset{S}{\otimes}R,\omega-F_i(\mathcal{X}_{i,1})}$, *where the polynomial ring R and the potential ω are determined by the boundary sequences except the sequence $\mathcal{X}_{i,1}$,*

$\mathcal{C}'\Big($  $\Big)_n \simeq \mathcal{C}'\Big($  $\Big)_n$

(3) *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{\mathbb{Q},0}$

$\mathcal{C}'\Big($  $\Big)_n$

$$= \overset{i}{\underset{j=1}{\boxtimes}} K\Big(L^{1;2}_{j,i}; x_{j,1}-x_{j,2}\Big)_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]}\Big|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}$$
$$\simeq \left(\,J_{F_i(\mathcal{X}_{i,1})} \to 0 \to J_{F_i(\mathcal{X}_{i,1})}\,\right)\{-in+i^2\}\,\langle i\rangle\,,$$

where $J_{F_i(\mathcal{X}_{i,1})}$ is Jacobi algebra for the polynomial $F_i(\mathcal{X}_{i,1})$), i.e.

$$J_{F_i(\mathcal{X}_{i,1})} = \mathbb{Q}[\mathcal{X}_{i,1}]\Big/\Big\langle \frac{\partial F_i}{\partial x_{1,1}},\cdots,\frac{\partial F_i}{\partial x_{i,1}}\Big\rangle\,.$$

Proof. **(1)** We describe the matrix factorization $\mathcal{C}'\Big($  $\Big)_n$ in $\mathrm{HMF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i,2}]\underset{S}{\otimes}R,\omega+F_i(\mathcal{X}_{i,2})}$ as (M_0,M_1,D_0,D_1), where M_j is an $R\underset{S}{\otimes}\mathbb{Q}[\mathcal{X}_{i,2}]$-module. Then we have

$\mathcal{C}'\Big($  $\Big)_n = (M_0,M_1,D_0,D_1)\overset{i}{\underset{j=1}{\boxtimes}} K\left(L^{1;2}_{j,i}; x_{j,1}-x_{j,2}\right)_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]}$.

We can regard this matrix factorization composed of $\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]\underset{S}{\otimes}R$-modules as a matrix factorization which consists of infinite rank $\mathbb{Q}[\mathcal{X}_{i,1}]\underset{S}{\otimes}R$-modules. Because the potential of this matrix factorization is in the polynomial ring $\mathbb{Q}[\mathcal{X}_{i,1}]\underset{S}{\otimes}R$. Moreover, the polynomials $x_{j,1}-x_{j,2}|_{\mathcal{X}_{i,1}=0}$ ($j=1,2,\cdots,i$) form regular in

$\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]\underset{S}{\otimes}R$. Thus, we can apply Corollary 2.15 to $(M_0,M_1,D_0,D_1)\boxtimes_{j=1}^{i} K\left(L^{1;2}_{j,i};x_{j,1}-x_{j,2}\right)_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]}$. Then we have

$$(M_0,M_1,D_0,D_1)\,|_{\mathcal{X}_{i,1}\to\mathcal{X}_{i,2}}\boxtimes\left(\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]\underset{S}{\otimes}R/\left\langle x_{1,1}-x_{1,2},\cdots,x_{i,1}-x_{i,2}\right\rangle,0,0,0\right)$$

The quotient polynomial ring $\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]\underset{S}{\otimes}R/\left\langle x_{1,1}-x_{1,2},\cdots,x_{i,1}-x_{i,2}\right\rangle$ is isomorphic to $\mathbb{Q}[\mathcal{X}_{i,1}]\underset{S}{\otimes}R$. Thus, we obtain the right-hand matrix factorization.

(2) This proof is similar to the proof of (1).

(3) We have

$$\mathop{\boxtimes}_{j=1}^{i} K\Big(L^{1;2}_{j,i};x_{j,1}-x_{j,2}\Big)_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]}\Big|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}=\mathop{\boxtimes}_{j=1}^{i}\Big(\mathbb{Q}[\mathcal{X}_{i,1}],\mathbb{Q}[\mathcal{X}_{i,1}]\{2j-1-n\},L^{1;2}_{j,i}|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}0\Big).$$

The polynomial $L^{1;2}_{j,i}|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}$ is

$$\frac{F_i(\cdots,x_{j-1,2},x_{j,1},x_{j+1,1},\cdots)-F_i(\cdots,x_{j-1,2},x_{j,2},x_{j+1,1},\cdots)}{x_{j,1}-x_{j,2}}\Big|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}=\frac{\partial F_i(\mathcal{X}_{i,1})}{\partial x_{j,1}}.$$

Hence, we apply Theorem 2.13 to these polynomials of the matrix factorization after using Proposition 2.5 and 2.8;

$$\begin{aligned}
&\mathop{\boxtimes}_{j=1}^{i} K\Big(L^{1;2}_{j,i};x_{j,1}-x_{j,2}\Big)_{\mathbb{Q}[\mathcal{X}_{i,1},\mathcal{X}_{i,2}]}\Big|_{\mathcal{X}_{i,2}\to\mathcal{X}_{i,1}}\\
\simeq\;&\mathop{\boxtimes}_{j=1}^{i}\left(\mathbb{Q}[\mathcal{X}_{i,1}],\mathbb{Q}[\mathcal{X}_{i,1}]\{n+1-2j\},0,\frac{\partial F_i(\mathcal{X}_{i,1})}{\partial x_{j,1}}\right)\{-in+i^2\}\left\langle i\right\rangle\\
\simeq\;&\left(J_{F_i(\mathcal{X}_{i,1})},0,0,0\right)\left\{-in+i^2\right\}\left\langle i\right\rangle.
\end{aligned}$$

□

A matrix factorization for the loop colored i, $\bigcirc i$, is defined to be the above matrix factorization;

$$\mathrm{MF}'\Big(\bigcirc i\Big):=\left(J_{F_i(\mathcal{X}_{i,1})},0,0,0\right)\left\{-in+i^2\right\}\left\langle i\right\rangle.$$

Corollary 3.6. *The Euler characteristic of the homology* $\mathrm{H}(\mathcal{C}'(\bigcirc i)_n)$ *equals MOY polynomial for the loop colored* i*;*

$$\overline{\chi}\Big(\mathrm{H}(\mathcal{C}'\Big(\bigcirc i\Big)_n)\Big)=\begin{bmatrix}n\\ i\end{bmatrix}.$$

Proof. By definition of Euler characteristic, the left-hand side of this equation equals the Poincaré series of Jacobi algebra times q^{-in+i^2}. Thus, we obtain the right-hand side of this equation by using the equation (7) in Section 3.1. □

Proposition 3.7. *Let* R *be a polynomial ring generated by variables of sequences* $\mathcal{X}_{i,1}$*,* $\mathcal{X}_{i,2}$*,* $\mathcal{X}_{i,3}$*,* $\mathcal{X}_{i,4}$ *and let* ω *be a polynomial*

$$F_{i_4}(x_{1,4},\cdots,x_{i_4,4})-F_{i_1}(x_{1,1},\cdots,x_{i_1,1})-F_{i_2}(x_{1,2},\cdots,x_{i_2,2})-F_{i_3}(x_{1,3},\cdots,x_{i_3,3}).$$

(1) *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{R,\omega}$



(2) *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{R,-\omega}$

$\mathcal{C}\Bigg($



$\Bigg)_n \simeq \mathcal{C}\Bigg($



$\Bigg)_n$,

where $1 \leq i_1, i_2, i_3 \leq n-2$, $i_5 = i_1 + i_2 \leq n-1$, $i_6 = i_2 + i_3 \leq n-1$ *and* $i_4 = i_1 + i_2 + i_3 \leq n$.

Proof. **(1)** We have

$\mathcal{C}\Bigg($



$\Bigg)_n$

$$= \boxtimes_{j=1}^{i_4} K\Big(\Lambda^{5;3,4}_{j,i_5;i_3}; x_{j,4} - X^{5;3}_{j,i_5;i_3}\Big)_{\mathbb{Q}[\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4},\mathcal{X}_{i_5,5}]} \boxtimes_{j=1}^{i_5} K\Big(\Lambda^{1;2,5}_{j,i_1;i_2}; x_{j,5} - X^{1;2}_{j,i_1;i_2}\Big)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_5,5}]}.$$

Since the potential of this matrix factorization does not include the variables of $\mathcal{X}_{i_5,5}$ and

$$(x_{1,5} - X^{1;2}_{1,i_1;i_2}, \cdots, x_{i_5,5} - X^{1;2}_{i_5,i_1;i_2})|_{(\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4})=(\underline{0})} = (\mathcal{X}_{i_5,5})$$

is obviously a regular sequence, we can apply Corollary 2.15 to these variables. Then the matrix factorization is isomorphic to

$$\boxtimes_{j=1}^{i_4} K\Big(\Lambda^{5;3,4}_{j,i_5;i_3}; x_{j,4} - X^{5;3}_{j,i_5;i_3}\Big)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4},\mathcal{X}_{i_5,5}]/\left\langle x_{1,5}-X^{1;2}_{1,i_1;i_2},\cdots,x_{i_5,5}-X^{1;2}_{i_5,i_1;i_2}\right\rangle}.$$

In the quotient ring $\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4},\mathcal{X}_{i_5,5}]/\left\langle x_{1,5} - X^{1;2}_{1,i_1;i_2}, \cdots, x_{i_5,5} - X^{1;2}_{i_5,i_1;i_2}\right\rangle$, the polynomial $X^{5;3}_{j,i_5;i_3}$ equals the $2j$-grading term of

$$(1 + x_{1,1} + x_{2,1} + \cdots + x_{i_1,1})(1 + x_{1,2} + x_{2,2} + \cdots + x_{i_2,2})(1 + x_{1,3} + x_{2,3} + \cdots + x_{i_3,3}).$$

If $X^{1;2;3}_{j,i_1;i_2;i_3}$ denotes the $2j$-grading term of

$$(1 + x_{1,1} + x_{2,1} + \cdots + x_{i_1,1})(1 + x_{1,2} + x_{2,2} + \cdots + x_{i_2,2})(1 + x_{1,3} + x_{2,3} + \cdots + x_{i_3,3}),$$

the polynomial $\Lambda^{5;3,4}_{j,i_5;i_3}$ equals to

$$\frac{F_{i_4}(\cdots, X^{1;2;3}_{j-1,i_1;i_2;i_3}, x_{j,4}, x_{j+1,4}, \cdots) - F_{i_4}(\cdots, X^{1;2;3}_{j-1,i_1;i_2;i_3}, X^{1;2;3}_{j,i_1;i_2;i_3}, x_{j+1,4}, \cdots)}{x_{j,4} - X^{1;2;3}_{j,i_1;i_2;i_3}}.$$

Let $\Lambda^{1;2;3,4}_{j,i_1;i_2;i_3}$ denote this polynomial. Then the matrix factorization is isomorphic to

$$\boxtimes_{j=1}^{i_4} K\Big(\Lambda^{1;2;3,4}_{j,i_1;i_2;i_3}; x_{j,4} - X^{1;2;3}_{j,i_1;i_2;i_3}\Big)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4}]}.$$

The other one is

$\mathcal{C}\Bigg($



$\Bigg)_n$

$$= \boxtimes_{j=1}^{i_4} K\Big(\Lambda^{1;6,4}_{j,i_1;i_6}; x_{j,4} - X^{1;6}_{j,i_1;i_6}\Big)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_4,4},\mathcal{X}_{i_6,6}]} \boxtimes_{j=1}^{i_6} K\Big(\Lambda^{2;3,6}_{j,i_2;i_3}; x_{j,6} - X^{2;3}_{j,i_2;i_3}\Big)_{\mathbb{Q}[\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_6,6}]}$$

Since the potential of this matrix factorization does not include the variables of $\mathcal{X}_{i_6,6}$ and

$$(x_{1,6}-X^{2;3}_{1,i_2;i_3},\cdots,x_{i_6,6}-X^{2;3}_{i_6,i_2;i_4})|_{(\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4})=(\underline{0})}=(\mathcal{X}_{i_6,6})$$

is a regular sequence in $\mathbb{Q}[\mathcal{X}_{i_1,1}\mathcal{X}_{i_2,2}\mathcal{X}_{i_3,3}\mathcal{X}_{i_4,4}\mathcal{X}_{i_6,6}]$, we can apply Corollary 2.15 to these variables. We similarly obtain the result that the matrix factorization is isomorphic to

$$\mathop{\boxtimes}_{j=1}^{i_4} K\Big(\Lambda^{1;2;3,4}_{j,i_1;i_2;i_3};x_{j,4}-X^{1;2;3}_{j,i_1;i_2;i_3}\Big)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4}]}.$$

(2) The proof of this proposition is similar to (1). By using Corollary 2.15, we find that the left-hand and right-hand matrix factorization are isomorphic to

$$\mathop{\boxtimes}_{j=1}^{i_4} K\Big(V^{1;2;3,4}_{j,i_1;i_2;i_3};X^{1;2;3}_{j,i_1;i_2;i_3}-x_{j,4}\Big)_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_2,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_4,4}]}\{-i_1i_2-i_2i_3-i_1i_3\},$$

where the polynomial $V^{1;2;3,4}_{j,i_1;i_2;i_3}$ is

$$\frac{F_{i_4}(\cdots,x_{j-1,4},X^{1;2;3}_{j,i_1;i_2;i_3},X^{1;2;3}_{j+1,i_1;i_2;i_3},\cdots)-F_{i_4}(\cdots,x_{j-1,4},x_{j,4},X^{1;2;3}_{j+1,i_1;i_2;i_3},\cdots)}{X^{1;2;3}_{j,i_1;i_2;i_3}-x_{j,4}}.$$

□

For $0\le n_2\le j\le n_1$, we define the positive integer $p(j,n_1;n_2)$ by

$$\begin{bmatrix} n_1 \\ n_2 \end{bmatrix}=\sum_{j=0}^{n_1n_2-n_2^2}p(j,n_1;n_2)\;q^{-n_1n_2+n_2^2+2j}.$$

Proposition 3.8. **(1)** *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2}],F_{i_3}(\mathcal{X}_{i_3,1})-F_{i_3}(\mathcal{X}_{i_3,2})}$

$\mathcal{C}\Bigg($



$\Bigg)_n \simeq \bigoplus_{j=0}^{i_1i_3-i_1^2}\Bigg(\mathcal{C}\Bigg($



$\Bigg)_n\{-i_1i_3+i_1^2+2j\}\Bigg)^{\oplus p(j,i_3,i_1)},$

(2) *There is the following isomorphism in* $\mathrm{HMF}^{gr}_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2}],F_{i_1}(\mathcal{X}_{i_1,1})-F_{i_1}(\mathcal{X}_{i_1,2})}$

$\mathcal{C}\Bigg($



$\Bigg)_n \simeq \bigoplus_{j=0}^{i_2(n-i_1)-i_2^2}\Bigg(\mathcal{C}\Bigg($



$\Bigg)_n\{-i_2(n-i_1)+i_2^2+2j\}\Bigg)^{\oplus p(j,n-i_1,i_2)}\langle i_2\rangle,$

where $1\le i_1,i_2\le n-1$ *and* $i_3=i_1+i_2\le n$.

Proof. **(1)** We have

$\mathcal{C}\Bigg($



$\Bigg)_n$

$$=\mathop{\boxtimes}_{j=1}^{i_3} K\Big(\Lambda^{3;4,1}_{j,i_1;i_2};x_{j,1}+X^{3;4}_{j,i_1;i_2}\Big)_{\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_1,3},\mathcal{X}_{i_2,4}]}\mathop{\boxtimes}_{j=1}^{i_3} K\Big(V^{3;4,2}_{j,i_1;i_2};x_{j,2}+X^{3;4}_{j,i_1;i_2}\Big)_{\mathbb{Q}[\mathcal{X}_{i_3,2},\mathcal{X}_{i_1,3},\mathcal{X}_{i_2,4}]}\{-i_1i_2\}.$$

The potential of this matrix factorization does not include the variables of $\mathcal{X}_{i_1,3}$, $\mathcal{X}_{i_2,4}$ and the sequence

$$(x_{1,2}+X^{3;4}_{1,i_1;i_2},x_{2,2}+X^{3;4}_{2,i_1;i_2},\cdots,x_{i_3,2}+X^{3;4}_{i_3,i_1;i_2})\Big|_{(\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2})=(\underline{0})}=(X^{3;4}_{1,i_1;i_2},X^{3;4}_{2,i_1;i_2},\cdots,X^{3;4}_{i_3,i_1;i_2})$$

is regular by Proposition 3.1, we can apply Corollary 2.15 to the matrix factorization. Thus we have

$$\boxtimes_{j=1}^{i_3} K\left(\Lambda^{3;4,1}_{j,i_1;i_2}; x_{j,1}-x_{j,2}\right)_{\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2},\mathcal{X}_{i_1,3},\mathcal{X}_{i_2,4}]/\left\langle x_{1,2}+X^{3;4}_{1,i_1;i_2},\cdots,x_{i_3,2}+X^{3;4}_{i_3,i_1;i_2}\right\rangle}\{-i_1i_2\}.$$

The polynomial $\Lambda^{3;4,1}_{j,i_1;i_2}$ is equal to L^{12}_{j,i_3} in the quotient ring

$$\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2},\mathcal{X}_{i_1,3},\mathcal{X}_{i_2,4}]/\left\langle x_{1,2}+X^{3;4}_{1,i_1;i_2},\cdots,x_{i_3,2}+X^{3;4}_{i_3,i_1;i_2}\right\rangle.$$

By the equation (4) in Section 3.1, the Poincaré series of the quotient ring is

$$\begin{aligned}&P_q(\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2},\mathcal{X}_{i_1,3},\mathcal{X}_{i_2,4}]/\left\langle x_{1,2}+X^{3;4}_{1,i_1;i_2},\cdots,x_{i_3,2}+X^{3;4}_{i_3,i_1;i_2}\right\rangle)\\ = \;&\frac{(1-q^2)(1-q^4)\cdots(1-q^{2i_1+2i_2})}{(1-q^2)(1-q^4)\cdots(1-q^{2i_1})(1-q^2)(1-q^4)\cdots(1-q^{2i_2})}P_q(\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2}]).\end{aligned}$$

Since

$$\frac{(1-q^2)(1-q^4)\cdots(1-q^{2i_1+2i_2})}{(1-q^2)(1-q^4)\cdots(1-q^{2i_1})(1-q^2)(1-q^4)\cdots(1-q^{2i_2})}=q^{i_1i_2}\sum_{j=1}^{i_1i_3-i_1^2}p(j,i_3;i_1)\;q^{-i_1i_3+i_1^2+2j},$$

the quotient ring $\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2},\mathcal{X}_{i_1,3},\mathcal{X}_{i_2,4}]/\left\langle x_{1,2}+X^{3;4}_{1,i_1;i_2},\cdots,x_{i_3,2}+X^{3;4}_{i_3,i_1;i_2}\right\rangle\{-i_1i_2\}$ is isomorphic to

$$\bigoplus_{j=0}^{i_1i_3-i_1^2}\left(\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2}]\left\{-i_1i_3+i_1^2+2j\right\}\right)^{\oplus p(j,i_3;i_1)}$$

as a $\mathbb{Z}$-graded $\mathbb{Q}[\mathcal{X}_{i_3,1},\mathcal{X}_{i_3,2}]$-algebra. Thus, we obtain the matrix factorization

$$\bigoplus_{j=0}^{i_1i_3-i_1^2}\left(\mathcal{C}\left(\begin{array}{c}\mathcal{X}_{i_3,1}\\ \uparrow{\scriptstyle i_3}\\ \mathcal{X}_{i_3,2}\end{array}\right)_n\{-i_1i_3+i_1^2+2j\}\right)^{\oplus p(j,i_3,i_1)}.$$

(2) We have

$\mathcal{C}\Bigg($



$\Bigg)_n$

$$= \boxtimes_{j=1}^{i_3} K(V^{2;4,3}_{j,i_1;i_2}; x_{j,3}-X^{2;4}_{j,i_1;i_2})_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_2,4}]}\{-i_1i_2\}\boxtimes_{j=1}^{i_3} K(\Lambda^{1;4,3}_{j,i_1;i_2}; X^{1;4}_{j,i_1;i_2}-x_{j,3})_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_2,4}]}$$

The potential of this matrix factorization does not include the variables of $\mathcal{X}_{i_3,3}$, $\mathcal{X}_{i_2,4}$ and the sequence

$$(X^{1;4}_{1,i_1;i_2}-x_{1,3},X^{1;4}_{2,i_1;i_2}-x_{2,3},\cdots,X^{1;4}_{i_3,i_1;i_2}-x_{i_3,3})\Big|_{(\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2})=(\underline{0})}=(x_{1,4}-x_{1,3},\cdots,x_{i_2,4}-x_{i_2,3},-x_{i_2+1,3},\cdots,-x_{i_3,3})$$

is regular in $\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_2,4}]$. Thus we can apply Corollary 2.15 to the matrix factorization. Then we have

$$\begin{aligned}&\boxtimes_{j=1}^{i_3} K(\Lambda^{2;4,3}_{j,i_1;i_2}; X^{1;4}_{j,i_1;i_2}-X^{2;4}_{j,i_1;i_2})_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_3,3},\mathcal{X}_{i_2,4}]/\left\langle X^{1;4}_{1,i_1;i_2}-x_{1,3},\cdots,X^{1;4}_{i_3,i_1;i_2}-x_{i_3,3}\right\rangle}\{-i_1i_2\}\\ (*)\quad\simeq\;&\boxtimes_{j=1}^{i_3} K(\widetilde{\Lambda^{2;4,3}_{j,i_1;i_2}}; X^{1;4}_{j,i_1;i_2}-X^{2;4}_{j,i_1;i_2})_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]}\{-i_1i_2\},\end{aligned}$$

where

$$\widetilde{\Lambda^{2;4,3}_{j,i_1;i_2}}=\frac{F_{i_3}(X^{2;4}_{1,i_1;i_2},\cdots,X^{2;4}_{j-1,i_1;i_2},X^{1;4}_{j,i_1;i_2},\cdots,X^{1;4}_{i_3,i_1;i_2})-F_{i_3}(X^{2;4}_{1,i_1;i_2},\cdots,X^{2;4}_{j,i_1;i_2},X^{1;4}_{j+1,i_1;i_2},\cdots,X^{1;4}_{i_3,i_1;i_2})}{X^{1;4}_{j,i_1;i_2}-X^{2;4}_{j,i_1;i_2}}.$$

Since $X^{1;4}_{j,i_1;i_2} - X^{2;4}_{j,i_1;i_2}$ is the $2j$-grading term of

$$((x_{1,1}-x_{1,2})+(x_{2,1}-x_{2,2})+\cdots+(x_{i_1,1}-x_{i_1,2}))(1+x_{1,4}+x_{2,4}+\cdots+x_{i_2,4}),$$

the polynomials $X^{1;4}_{j,i_1;i_2} - X^{2;4}_{j,i_1;i_2}$ $(i_1+1 \le j \le i_3)$ can be described as the linear sum of the polynomials $X^{1;4}_{j,i_1;i_2} - X^{2;4}_{j,i_1;i_2}$ $(1 \le j \le i_1)$. Using Proposition 2.9 the matrix factorization $(*)$ is isomorphic to

$$\begin{aligned}&\mathop{\boxtimes}_{j=1}^{i_1} K(*; x_{j,1}-x_{j,2})_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]} \boxtimes \\ &\mathop{\boxtimes}_{k=i_1+1}^{i_3} (Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}], Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]\{2k-n-1\}, \widetilde{\Lambda^{2;4,3}_{k,i_1;i_2}}, 0)\,\{-i_1 i_2\}\,.\end{aligned}$$

Since the potential of the partial matrix factorization $\mathop{\boxtimes}_{j=1}^{i_1} K(*; x_{j,1}-x_{j,2})_{\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]}$ equals $F_{i_1}(\mathcal{X}_{i_1,1}) - F_{i_1}(\mathcal{X}_{i_1,2})$, by Theorem 2.10 the partial matrix factorization is isomorphic to

$\mathcal{C}\Bigg($  $\Bigg)_n \boxtimes (\mathbb{Q}[\mathcal{X}_{i_2,4}],0,0,0).$

The other partial matrix factorization $\boxtimes_{k=i_1+1}^{i_3}(Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}], Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]\{2k-n-1\}, \widetilde{\Lambda^{2;4,3}_{k,i_1;i_2}}, 0)\,\{-i_1 i_2\}$ is isomorphic to

$$\mathop{\boxtimes}_{k=i_1+1}^{i_3} (0, Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]/\left\langle \widetilde{\Lambda^{2;4,3}_{k,i_1;i_2}} \right\rangle \{2k-n-1\}, 0, 0).$$

Thus, we have

$\mathcal{C}\Bigg($  $\Bigg)_n \boxtimes \mathop{\boxtimes}_{k=i_1+1}^{i_3} (0, Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]/\left\langle \widetilde{\Lambda^{2;4,3}_{k,i_1;i_2}} \right\rangle \{2k-n-1\}, 0, 0)\,\{-i_1 i_2\}\,.$

Since we find that $\widetilde{\Lambda^{2;4,3}_{k,i_1;i_2}}\Big|_{(\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2})=(\underline{0})} = \frac{\partial F_{i_3}}{\partial x_j}(x_{1,4},\cdots,x_{i_1,4},0,\cdots,0)$ $\quad(i_1+1 \le j \le i_3)$ form regular sequence, these polynomials $\widetilde{\Lambda^{2;4,3}_{i_1+1,i_1;i_2}},\cdots,\widetilde{\Lambda^{2;4,3}_{i_3,i_1;i_2}}$ also form regular. The Poincaré series of the $\mathbb{Z}$-graded quotient ring

$$Q[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2},\mathcal{X}_{i_2,4}]/\left\langle \widetilde{\Lambda^{2;4,3}_{i_1+1,i_1;i_2}},\cdots,\widetilde{\Lambda^{2;4,3}_{i_3,i_1;i_2}} \right\rangle \{\sum_{k=i_1+1}^{i_3} 2k-n-1\}\,\{-i_1 i_2\}$$

equals to

$$\begin{aligned}&\frac{(1-q^{2(n-i_1)})(1-q^{2(n-i_1-1)})\cdots(1-q^{2(n-i_3+1)})}{(1-q^2)(1-q^4)\cdots(1-q^{2i_2})} P_q(\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2}]) \prod_{k=i_1+1}^{i_3} q^{2k-n-1} \times q^{-i_1 i_2} \\ = \;& \begin{bmatrix} n-i_1 \\ i_2 \end{bmatrix} P_q(\mathbb{Q}[\mathcal{X}_{i_1,1},\mathcal{X}_{i_1,2}]).\end{aligned}$$

Hence, the matrix factorization $\mathcal{C}\Bigg($  $\Bigg)_n$ is isomorphic to

$$\bigoplus_{j=0}^{i_2(n-i_1)-i_2^2}\left(\mathcal{C}'\left(\begin{array}{c}\mathcal{X}_{i_1,1}\\ \uparrow{\scriptstyle i_1}\\ \mathcal{X}_{i_1,2}\end{array}\right)_n\{-i_2(n-i_1)+i_2^2+2j\}\right)^{\oplus p(j,n-i_1,i_2)}\langle i_2\rangle\,.$$

□

Proposition 3.9. *There is the following isomorphism in* $\mathrm{HMF}_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4}],F_1(\mathcal{X}_{1,1})+F_j(\mathcal{X}_{j,2})-F_1(\mathcal{X}_{1,3})-F_j(\mathcal{X}_{j,4})}$

$(\mathbf{1})\mathcal{C}'\Bigg($



$\Bigg)_n \simeq \mathcal{C}'\Bigg($



$\Bigg)_n \bigoplus_{i=1}^{j-1}\mathcal{C}'\Bigg($



$\Bigg)_n \{2i-j\}$

$(\mathbf{2})\mathcal{C}'\Bigg($



$\Bigg)_n \simeq \mathcal{C}'\Bigg($



$\Bigg)_n \bigoplus_{k=1}^{n-j-1}\mathcal{C}'\Bigg($



$\Bigg)_n \{2k+j-n\}\langle 1\rangle\,.$

Proof. **(1)**We have



$$
\begin{aligned}
= \;& K\left(\begin{pmatrix} V^{1;6,5}_{1,1;1}\\ V^{1;6,5}_{2,1;1}\end{pmatrix};\begin{pmatrix} x_{1,1}+x_{1,6}-x_{1,5}\\ x_{1,1}x_{1,6}-x_{2,5}\end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{1,6},\mathcal{X}_{2,5}]}\\
&\boxtimes K\left(\begin{pmatrix} \Lambda^{3;8,5}_{1,1;1}\\ \Lambda^{3;8,5}_{2,1;1}\end{pmatrix};\begin{pmatrix} x_{1,5}-x_{1,3}-x_{1,8}\\ x_{2,5}-x_{1,3}x_{1,8}\end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,3},\mathcal{X}_{1,8},\mathcal{X}_{2,5}]}\{-1\}\\
&\boxtimes K\left(\begin{pmatrix} \Lambda^{6;7,2}_{1,1;j-1}\\ \Lambda^{6;7,2}_{2,1;j-1}\\ \Lambda^{6;7,2}_{3,1;j-1}\\ \vdots\\ \Lambda^{6;7,2}_{j-1,1;j-1}\\ \Lambda^{6;7,2}_{j,1;j-1}\end{pmatrix};\begin{pmatrix} x_{1,2}-x_{1,6}-x_{1,7}\\ x_{2,2}-x_{1,6}x_{1,7}-x_{2,7}\\ x_{3,2}-x_{1,6}x_{2,7}-x_{3,7}\\ \vdots\\ x_{j-1,2}-x_{1,6}x_{j-2,7}-x_{j-1,7}\\ x_{j,2}-x_{1,6}x_{j-1,7}\end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{j,2},\mathcal{X}_{1,6},\mathcal{X}_{j-1,7}]}\{-j+1\}\\
&\boxtimes K\left(\begin{pmatrix} V^{8;7,4}_{1,1;j-1}\\ V^{8;7,4}_{2,1;j-1}\\ V^{8;7,4}_{3,1;j-1}\\ \vdots\\ V^{8;7,4}_{j-1,1;j-1}\\ V^{8;7,4}_{j,1;j-1}\end{pmatrix};\begin{pmatrix} x_{1,7}+x_{1,8}-x_{1,4}\\ x_{1,7}x_{1,8}+x_{2,7}-x_{2,4}\\ x_{2,7}x_{1,8}+x_{3,7}-x_{3,4}\\ \vdots\\ x_{j-2,7}x_{1,8}+x_{j-1,7}-x_{j-1,4}\\ x_{j-1,7}x_{1,8}-x_{j,4}\end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,8},\mathcal{X}_{j-1,7},\mathcal{X}_{j,2}]}.
\end{aligned}
$$

We apply Corollary 2.15 to the matrix factorization. Then we obtain

$$
K\left(\begin{pmatrix} \Lambda^{3;8,5}_{2,1;1}\\ \Lambda^{6;7,2}_{j,1;j-1}\\ V^{8;7,4}_{1,1;j-1}\\ V^{8;7,4}_{2,1;j-1}\\ V^{8;7,4}_{3,1;j-1}\\ \vdots\\ V^{8;7,4}_{j-1,1;j-1}\\ V^{8;7,4}_{j,1;j-1}\end{pmatrix};\begin{pmatrix} x_{2,5}-x_{1,3}x_{1,8}\\ x_{j,2}-x_{1,6}x_{j-1,7}\\ x_{1,7}+x_{1,8}-x_{1,4}\\ x_{1,7}x_{1,8}+x_{2,7}-x_{2,4}\\ x_{2,7}x_{1,8}+x_{3,7}-x_{3,4}\\ \vdots\\ x_{j-2,7}x_{1,8}+x_{j-1,7}-x_{j-1,4}\\ x_{j-1,7}x_{1,8}-x_{j,4}\end{pmatrix}\right)_{R_1}\{-j\},
$$

where $R_1 = \mathbb{Q}\left[\begin{array}{c}\mathcal{X}_{1,1}, \mathcal{X}_{j,2}, \mathcal{X}_{1,3}, \mathcal{X}_{j,4},\\ \mathcal{X}_{2,5}, \mathcal{X}_{1,6}, \mathcal{X}_{j-1,7}, \mathcal{X}_{1,8}\end{array}\right] \Big/\Big\langle \begin{array}{c} x_{1,1}+x_{1,6}-x_{1,5}, x_{1,1}x_{1,6}-x_{2,5}, x_{1,5}-x_{1,3}-x_{1,8},\\ x_{1,2}-x_{1,6}-x_{1,7}, \cdots, x_{j,2}-x_{1,6}x_{j-1,7}\end{array}\Big\rangle$.
In the quotient ring, there are the following equations

$$\begin{aligned}
x_{1,5} &= x_{1,1}+x_{1,6},\\
x_{2,5} &= x_{1,1}x_{1,6},\\
x_{1,8} &= x_{1,1}-x_{1,3}+x_{1,6},\\
x_{k,7} &= \sum_{l=0}^{k}(-1)^l x_{1,6}^l x_{k-l,2}\,(=: A_k) \qquad (k=1,2,\cdots,j-1),
\end{aligned}$$

where $x_{0,2}=1$.
Then we can consider $R_1 \simeq \mathbb{Q}[\mathcal{X}_{1,1}, \mathcal{X}_{j,2}, \mathcal{X}_{1,3}, \mathcal{X}_{j,4}, \mathcal{X}_{1,6}]$. That is, the variables $x_{1,5}$, $x_{2,5}$, $x_{1,8}$ and $x_{k,7}$ can be removed from the quotient ring R_1 using the above equations. Therefore the matrix factorization is isomorphic to

$$K\left(\left(\begin{array}{c}
\widetilde{\Lambda_{2,1;1}^{3;8,5}}\\
\widetilde{\Lambda_{j,1;j-1}^{6;7,2}}\\
\widetilde{V_{1,1;j-1}^{8;7,4}}\\
\widetilde{V_{2,1;j-1}^{8;7,4}}\\
\vdots\\
\widetilde{V_{j-1,1;j-1}^{8;7,4}}\\
\widetilde{V_{j,1;j-1}^{8;7,4}}
\end{array}\right);\left(\begin{array}{c}
(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})\\
\sum_{l=0}^{j}(-1)^l x_{1,6}^l x_{j-l,2}\\
x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\
(x_{1,1}-x_{1,3})A_1+x_{2,2}-x_{2,4}\\
\vdots\\
(x_{1,1}-x_{1,3})A_{j-2}+x_{j-1,2}-x_{j-1,4}\\
(x_{1,1}-x_{1,3}+x_{1,6})A_{j-1}-x_{j,4}
\end{array}\right)\right)_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4},\mathcal{X}_{1,6}]}\{-j\} \qquad .$$

Theorem 2.13 is applied to $\sum_{l=0}^{j}(-1)^l x_{1,6}^l x_{j-l,2}$ of the matrix factorization. Then we obtain

$$K\left(\left(\begin{array}{c}
\widehat{\Lambda_{2,1;1}^{3;8,5}}\\
\widehat{V_{1,1;j-1}^{8;7,4}}\\
\widehat{V_{2,1;j-1}^{8;7,4}}\\
\vdots\\
\widehat{V_{j,1;j-1}^{8;7,4}}
\end{array}\right);\left(\begin{array}{c}
(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})\\
x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\
(x_{1,1}-x_{1,3})A_1+x_{2,2}-x_{2,4}\\
\vdots\\
(x_{1,1}-x_{1,3})A_{j-1}+x_{j,2}-x_{j,4}
\end{array}\right)\right)_{R_1{}'}\{-j\},$$

where $R_1{}' = \mathbb{Q}[\mathcal{X}_{1,1}, \mathcal{X}_{j,2}, \mathcal{X}_{1,3}, \mathcal{X}_{j,4}, \mathcal{X}_{1,6}] \Big/\Big\langle \sum_{l=0}^{j}(-1)^l x_{1,6}^l x_{j-l,2}\Big\rangle$, $\widehat{\Lambda_{2,1;1}^{3;8,5}} = \widetilde{\Lambda_{2,1;1}^{3;8,5}}$ and $\widehat{V_{i,1;j-1}^{8;7,4}} = \widetilde{V_{i,1;j-1}^{8;7,4}}$ in the quotient ring $R_1{}'$. Since the polynomials A_k are described as

$$\begin{aligned}
A_k &= \left(\sum_{l_1=0}^{k-1}(-1)^{k-l_1}x_{1,6}^{k-1-l_1}\left(\sum_{l_2=0}^{l_1}(-1)^{l_1-l_2}x_{1,3}^{l_1-l_2}x_{l_2,2}\right)\right)(x_{1,6}-x_{1,3})+\sum_{l_3=0}^{k}(-1)^{k-l_3}x_{1,3}^{k-l_3}x_{l_3,2}\\
&= u_{k-1}(x_{1,6}-x_1,3)+v_k,
\end{aligned}$$

the above matrix factorization is isomorphic to

$$K\left(\left(\begin{array}{c}\widehat{\Lambda^{3;8,5}_{2,1;1}}+\sum_{k=2}^{j}u_{k-2}\widehat{V^{8;7,4}_{k,1;j-1}}\\ \widehat{V^{8;7,4}_{1,1;j-1}}\\ \widehat{V^{8;7,4}_{2,1;j-1}}\\ \vdots\\ \widehat{V^{8;7,4}_{j,1;j-1}}\end{array}\right);\left(\begin{array}{c}(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})\\ x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\ (x_{1,1}-x_{1,3})v_1+x_{2,2}-x_{2,4}\\ \vdots\\ (x_{1,1}-x_{1,3})v_{j-1}+x_{j,2}-x_{j,4}\end{array}\right)\right)_{R_1'}\{-j\}.$$

We can apply Corollary 2.11 to the matrix factorization, and then there are polynomials $a\in R_1'$ and $a_k\in\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4}](:=R_1'')$ $(k=1,\cdots j)$ to give an isomorphism between the above matrix factorization and the following matrix factorization

$$K\left(\left(\begin{array}{c}a\\ a_1\\ a_2\\ \vdots\\ a_j\end{array}\right);\left(\begin{array}{c}(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})\\ x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\ (x_{1,1}-x_{1,3})v_1+x_{2,2}-x_{2,4}\\ \vdots\\ (x_{1,1}-x_{1,3})v_{j-1}+x_{j,2}-x_{j,4}\end{array}\right)\right)_{R_1'}\{-j\}. \tag{11}$$

The partial matrix factorization $K(a;(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3}))$ has the potential

$$a(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})=\omega-\sum_{i=1}^{j}a_i(-(x_{1,1}-x_{1,3})(\sum_{k=0}^{i-1}(-1)^{i-1-k}x_{1,3}^{i-1-k}x_{k,2})+x_{i,2}-x_{i,4})\in R_1''.$$

Thus we have

$$\begin{array}{cl} & k(a;(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3}))_{R_1'}\\ \simeq & \left(\left(\begin{array}{r}R_1''\\ (x_{1,6}-x_{1,3})R_1''\\ x_{1,6}(x_{1,6}-x_{1,3})R_1''\\ \vdots\\ x_{1,6}^{j-3}(x_{1,6}-x_{1,3})R_1''\\ x_{1,6}^{j-2}(x_{1,6}-x_{1,3})R_1''\end{array}\right),\left(\begin{array}{r}R_1''\{3-n\}\\ x_{1,6}R_1''\{3-n\}\\ x_{1,6}^{2}R_1''\{3-n\}\\ \vdots\\ x_{1,6}^{j-2}R_1''\{3-n\}\\ \alpha R_1''\{3-n\}\end{array}\right),f_0,f_1\right).\end{array}$$

where

$$\begin{array}{rcl} f_0 & = & \left(\begin{array}{cccc}0 & a(x_{1,6}-x_{1,3}) & & \\ \vdots & & \ddots & \\ 0 & & & a(x_{1,6}-x_{1,3})\\ \frac{a}{\alpha} & 0 & \cdots & 0\end{array}\right),\\ f_1 & = & \left(\begin{array}{cccc}0 & \cdots & 0 & \alpha(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})\\ x_{1,1}-x_{1,3} & & & 0\\ & \ddots & & \vdots\\ & & x_{1,1}-x_{1,3} & 0\end{array}\right),\\ \alpha & = & \sum_{k_1=0}^{j-1}(-1)^{j-1-k_1}x_{1,6}^{j-1-k_1}\left(\sum_{k_2=0}^{k_1}(-1)^{k_1-k_2}x_{1,3}^{k_1-k_2}x_{k_2,2}\right).\end{array}$$

Remark that it is obvious to find

$$\alpha(x_{1,6}-x_{1,3})=\sum_{k=0}^{j}(-1)^{j-k}x_{1,3}^{j-k}x_{k,2}.$$

Hence the partial matrix factorization splits into the following direct sum

$$
\begin{aligned}
&(\ (x_{1,6}-x_{1,3})R_1'' \xrightarrow{a(x_{1,6}-x_{1,3})} R_1''\{3-n\} \xrightarrow{x_{1,1}-x_{1,3}} (x_{1,6}-x_{1,3})R_1''\ )\\
&\oplus(\ x_{1,6}(x_{1,6}-x_{1,3})R_1'' \xrightarrow{a(x_{1,6}-x_{1,3})} x_{1,6}R_1''\{3-n\} \xrightarrow{x_{1,1}-x_{1,3}} x_{1,6}(x_{1,6}-x_{1,3})R_1''\ )\\
&\vdots\\
&\oplus(\ x_{1,6}^{j-2}(x_{1,6}-x_{1,3})R_1'' \xrightarrow{a(x_{1,6}-x_{1,3})} x_{1,6}^{j-2}R_1''\{3-n\} \xrightarrow{x_{1,1}-x_{1,3}} x_{1,6}^{j-2}(x_{1,6}-x_{1,3})R_1''\ )\\
&\oplus(\ R_1'' \xrightarrow{\frac{a}{\alpha}} \alpha R_1''\{3-n\} \xrightarrow{\alpha(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})} R_1''\ ),
\end{aligned}
$$

$$
\begin{aligned}
\simeq\ &\left.\begin{aligned}
&(\ R_1'' \xrightarrow{a(x_{1,6}-x_{1,3})} R_1''\{1-n\} \xrightarrow{x_{1,1}-x_{1,3}} R_1''\ )\{2\}\\
&\vdots\\
&\oplus(\ R_1'' \xrightarrow{a(x_{1,6}-x_{1,3})} R_1''\{1-n\} \xrightarrow{x_{1,1}-x_{1,3}} R_1''\ )\{2j-2\}\\
&\oplus(\ R_1'' \xrightarrow{\frac{a}{\alpha}} R_1''\{2j+1-n\} \xrightarrow{\alpha(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3})} R_1''\ ).
\end{aligned}\right.\\
\simeq\ &\bigoplus_{i=1}^{j-1} K(a(x_{1,6}-x_{1,3});x_{1,1}-x_{1,3})_{R_1''}\{2i\} \oplus K(\frac{a}{\alpha};\alpha(x_{1,6}-x_{1,3})(x_{1,1}-x_{1,3}))_{R_1''}
\end{aligned}
$$

By the decomposition and using Theorem 2.10, The total matrix factorization (11) is isomorphic to

$$
\begin{aligned}
&\bigoplus_{i=1}^{j-1} K\left(\left(\begin{array}{c} a(x_{1,6}-x_{1,3})\\ a_1\\ a_2\\ \vdots\\ a_j \end{array}\right);\left(\begin{array}{c} x_{1,1}-x_{1,3}\\ x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\ (x_{1,1}-x_{1,3})(\sum_{k=0}^{1}(-1)^{1-k}x_{1,3}^{1-k}x_{k,2})+x_{2,2}-x_{2,4}\\ \vdots\\ (x_{1,1}-x_{1,3})(\sum_{k=0}^{j-1}(-1)^{j-1-k}x_{1,3}^{j-1-k}x_{k,2})+x_{j,2}-x_{j,4} \end{array}\right)\right)_{R_1''}\{2i-j\}\\
&\bigoplus K\left(\left(\begin{array}{c} a_1\\ a_2\\ \vdots\\ a_j\\ \frac{a}{\alpha} \end{array}\right);\left(\begin{array}{c} x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\ (x_{1,1}-x_{1,3})(\sum_{k=0}^{1}(-1)^{1-k}x_{1,3}^{1-k}x_{k,2})+x_{2,2}-x_{2,4}\\ \vdots\\ (x_{1,1}-x_{1,3})(\sum_{k=0}^{j-1}(-1)^{j-1-k}x_{1,3}^{j-1-k}x_{k,2})+x_{j,2}-x_{j,4}\\ (x_{1,1}-x_{1,3})(\sum_{k=0}^{j}(-1)^{j-k}x_{1,3}^{j-k}x_{k,2}) \end{array}\right)\right)_{R_1''}\{-j\}\\
\simeq &\bigoplus_{i=1}^{j-1} K\left(\left(\begin{array}{c} a(x_{1,6}-x_{1,3})+\sum_{l=1}^{j}(\sum_{k=0}^{l-1}(-1)^{l-2-k}x_{1,3}^{l-1-k}x_{k,2})a_l\\ a_1\\ a_2\\ \vdots\\ a_j \end{array}\right);\left(\begin{array}{c} x_{1,1}-x_{1,3}\\ x_{1,2}-x_{1,4}\\ x_{2,2}-x_{2,4}\\ \vdots\\ x_{j,2}-x_{j,4} \end{array}\right)\right)_{R_1''}\{2i-j\}\\
&\bigoplus K\left(\left(\begin{array}{c} \sum_{k=1}^{j+1}(-1)^{k-1}a_k x_{1,3}^{k-1}\\ \sum_{k=2}^{j+1}(-1)^{k-2}a_k x_{1,3}^{k-2}\\ \vdots\\ a_{j-1}-x_{1,3}a_j+x_{1,3}^2a_{j+1}\\ a_j-x_{1,3}a_{j+1}\\ a_{j+1} \end{array}\right);\left(\begin{array}{c} x_{1,1}+x_{1,2}-x_{1,3}-x_{1,4}\\ x_{1,1}x_{1,2}-x_{1,3}x_{1,4}+x_{2,2}-x_{2,4}\\ \vdots\\ x_{1,1}x_{j-2,2}-x_{1,3}x_{j-2,4}+x_{j-1,2}-x_{j-1,4}\\ x_{1,1}x_{j-1,2}-x_{1,3}x_{j-1,4}+x_{j,2}-x_{j,4}\\ x_{1,1}x_{j,2}-x_{1,3}x_{j,4} \end{array}\right)\right)_{R_1''}\{-j\}
\end{aligned}
$$

where $a_{j+1} = \frac{a}{\alpha}$.
Using Theorem 2.10, we find the above matrix factorization is isomorphic to



(2) We have

$$K\left(\begin{pmatrix} \Lambda^{1;5,6}_{1,1;j} \\ \Lambda^{1;5,6}_{2,1;j} \\ \vdots \\ \Lambda^{1;5,6}_{j,1;j} \\ \Lambda^{1;5,6}_{j+1,1;j} \end{pmatrix} ; \begin{pmatrix} x_{1,6} - x_{1,1} - x_{1,5} \\ x_{2,6} - x_{1,1}x_{1,5} - x_{2,5} \\ \vdots \\ x_{j,6} - x_{1,1}x_{j-1,5} - x_{j,5} \\ x_{j+1,6} - x_{1,1}x_{j,5} - x_{j+1,5} \end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,1}\mathcal{X}_{j,5}\mathcal{X}_{j+1,6}]} \{-j\}$$

$$\boxtimes K\left(\begin{pmatrix} V^{3;5,8}_{1,1;j} \\ V^{3;5,8}_{2,1;j} \\ \vdots \\ V^{3;5,8}_{j,1;j} \\ V^{3;5,8}_{j+1,1;j} \end{pmatrix} ; \begin{pmatrix} x_{1,3} + x_{1,5} - x_{1,8} \\ x_{1,3}x_{1,5} + x_{2,5} - x_{2,8} \\ \vdots \\ x_{1,3}x_{j-1,5} + x_{j,5} - x_{j,8} \\ x_{1,3}x_{j,5} - x_{j+1,8} \end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,3}\mathcal{X}_{j,5}\mathcal{X}_{j+1,8}]}$$

$$\boxtimes K\left(\begin{pmatrix} V^{7;2,6}_{1,1;j} \\ V^{7;2,6}_{2,1;j} \\ \vdots \\ V^{7;2,6}_{j,1;j} \\ V^{7;2,6}_{j+1,1;j} \end{pmatrix} ; \begin{pmatrix} x_{1,7} + x_{1,2} - x_{1,6} \\ x_{1,7}x_{1,2} + x_{2,2} - x_{2,6} \\ \vdots \\ x_{1,7}x_{j-1,2} + x_{j,2} - x_{j,6} \\ x_{1,7}x_{j,2} - x_{j+1,6} \end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,7}\mathcal{X}_{j,2}\mathcal{X}_{j+1,6}]}$$

$$\boxtimes K\left(\begin{pmatrix} \Lambda^{7;4,8}_{1,1;j} \\ \Lambda^{7;4,8}_{2,1;j} \\ \vdots \\ \Lambda^{7;4,8}_{j,1;j} \\ \Lambda^{7;4,8}_{j+1,1;j} \end{pmatrix} ; \begin{pmatrix} x_{1,8} - x_{1,7} - x_{1,4} \\ x_{2,8} - x_{1,7}x_{1,4} - x_{2,4} \\ \vdots \\ x_{j,8} - x_{1,7}x_{j-1,4} - x_{j,4} \\ x_{j+1,8} - x_{1,7}x_{j,4} \end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,7}\mathcal{X}_{j,4}\mathcal{X}_{j+1,8}]} \{-j\}.$$

We apply Corollary 2.15 to the matrix factorization. Then we obtain

$$K\left(\begin{pmatrix} \Lambda^{1;5,6}_{1,1;j} \\ \vdots \\ \Lambda^{1;5,6}_{j+1,1;j} \\ V^{3;5,8}_{j+1,1;j} \end{pmatrix} ; \begin{pmatrix} x_{1,6} - x_{1,1} - x_{1,5} \\ \vdots \\ x_{j+1,6} - x_{1,1}x_{j,5} \\ x_{1,3}x_{j,5} - x_{j+1,8} \end{pmatrix}\right)_{R_2} \{-2j\},$$

where $R_2 = \mathbb{Q}\left[\begin{array}{c} \mathcal{X}_{1,1}, \mathcal{X}_{j,2}, \mathcal{X}_{1,3}, \mathcal{X}_{j,4}, \\ \mathcal{X}_{j,5}, \mathcal{X}_{j+1,6}, \mathcal{X}_{1,7}, \mathcal{X}_{j+1,8} \end{array}\right] \Big/ \left\langle \begin{array}{c} x_{1,3}+x_{1,5}-x_{1,8}, \cdots, x_{1,3}x_{j-1,5}+x_{j,5}-x_{j,8}, \\ x_{1,7}+x_{1,2}-x_{1,6}, \cdots, x_{1,7}x_{j,2}-x_{j+1,6}, \\ x_{1,8}-x_{1,7}-x_{1,4}, \cdots, x_{j+1,8}-x_{1,7}x_{j,4} \end{array}\right\rangle$

In the quotient ring, there are equations

$$\begin{array}{rcl} x_{k,5} &=& \left(\sum_{l=0}^{k-1}(-1)^{k-1-l}x_{1,3}^{k-1-l}x_{l,4}\right)x_{1,7} + \sum_{l=0}^{k}(-1)^{k-l}x_{1,3}^{k-l}x_{l,4} (=: B_k) \qquad (k=1,\cdots,j), \\ x_{1,6} &=& x_{1,7}+x_{1,2}, \\ && \vdots \\ x_{j+1,6} &=& x_{1,7}x_{j,2}, \\ x_{1,8} &=& x_{1,7}+x_{1,4}, \\ && \vdots \\ x_{j+1,8} &=& x_{1,7}x_{j,4}. \end{array}$$

Using the above equations, we find the matrix factorization is isomorphic to

$$\begin{array}{rl} & K\left(\left(\begin{array}{c} \widetilde{\Lambda^{1;5,6}_{1,1;j}} \\ \widetilde{\Lambda^{1;5,6}_{2,1;j}} \\ \vdots \\ \widetilde{\Lambda^{1;5,6}_{j,1;j}} \\ \widetilde{\Lambda^{1;5,6}_{j+1,1;j}} \\ \widetilde{V^{3;5,8}_{j+1,1;j}} \end{array}\right);\left(\begin{array}{c} s_1 \\ s_1x_{1,7}+s_2 \\ \vdots \\ s_{j-1}x_{1,7}+s_j \\ x_{1,7}x_{j,2}-x_{1,1}B_j \\ x_{1,3}B_j - x_{1,7}x_{j,4} \end{array}\right)\right)_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4},\mathcal{X}_{1,7}]} \qquad \{-2j\}, \\ \simeq & K\left(\left(\begin{array}{c} \widetilde{\Lambda^{1;5,6}_{1,1;j}} + x_{1,7}\widetilde{\Lambda^{1;5,6}_{2,1;j}} \\ \vdots \\ \widetilde{\Lambda^{1;5,6}_{j-1,1;j}} + x_{1,7}\widetilde{\Lambda^{1;5,6}_{j,1;j}} \\ \widetilde{\Lambda^{1;5,6}_{j,1;j}} \\ \widetilde{\Lambda^{1;5,6}_{j+1,1;j}} \\ \widetilde{V^{3;5,8}_{j+1,1;j}} \end{array}\right);\left(\begin{array}{c} s_1 \\ \vdots \\ s_{j-1} \\ s_j \\ x_{1,7}x_{j,2}-x_{1,1}B_j \\ (x_{1,3}-x_{1,7})(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}) \end{array}\right)\right)_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4},\mathcal{X}_{1,7}]} \qquad \{-2j\}, \end{array}$$

where $s_k = x_{k,2} + \sum_{l=0}^{k}(-1)^{k+1-l}x_{1,3}^{k-l}X^{1;4}_{l,1;j}$ and $\widetilde{\Lambda^{1;5,6}_{k,1;j}}$, $\widetilde{V^{3;5,8}_{j+1,1;j}}$ are derived from $\Lambda^{1;5,6}_{k,1;j}$, $V^{3;5,8}_{j+1,1;j}$ using the equations.

The polynomial $\widetilde{\Lambda^{1;5,6}_{j+1,1;j}}$ can be described by

$$\begin{array}{rcl} \widetilde{\Lambda^{1;5,6}_{j+1,1;j}} &=& \dfrac{F_{j+1}(X^{1;5}_{1,1;j},\cdots,X^{1;5}_{j,1;j},x_{j+1,6}) - F_{j+1}(X^{1;5}_{1,1;j},\cdots,X^{1;5}_{j,1;j},X^{1;5}_{j+1,1;j})}{x_{j+1,6}-X^{1;5}_{j+1,1;j}} \\ &=& c_0\left(X^{1;5}_{1,1;j}\right)^{n-j} + \cdots \\ &=& c_0(x_{1,7}+x_{1,1}-x_{1,3}+x_{1,4})^{n-j} + \cdots \\ &=& c_0x_{1,7}^{n-j} + c_1x_{1,7}^{n-j-1} + \cdots + c_{n-j}, \end{array}$$

where $c_0 \in \mathbb{Q}$ and $c_k \in \mathbb{Q}[\mathcal{X}_{1,1}, \mathcal{X}_{j,2}, \mathcal{X}_{1,3}, \mathcal{X}_{j,4}](:= R_2'')$ $(k = 1, \cdots, n-j)$. Using Theorem 2.13, we have

$$
\begin{aligned}
& K\left(\left(\begin{array}{c}
\widetilde{\Lambda^{1;5,6}_{1,1;j}} + x_{1,7}\widetilde{\Lambda^{1;5,6}_{2,1;j}}\\
\vdots\\
\widetilde{\Lambda^{1;5,6}_{j-1,1;j}} + x_{1,7}\widetilde{\Lambda^{1;5,6}_{j,1;j}}\\
\widetilde{\Lambda^{1;5,6}_{j,1;j}}\\
x_{1,7}x_{j,2} - x_{1,1}B_j\\
\widetilde{V^{3;5,8}_{j+1,1;j}}
\end{array}\right);\left(\begin{array}{c}
s_1\\
\vdots\\
s_{j-1}\\
s_j\\
\widetilde{\Lambda^{1;5,6}_{j+1,1;j}}\\
(x_{1,3}-x_{1,7})(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4})
\end{array}\right)\right)_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4},\mathcal{X}_{1,7}]} \{-2j\}\{2j+1-n\}\left\langle 1\right\rangle,\\
\simeq\ & K\left(\left(\begin{array}{c}
\widetilde{\Lambda^{1;5,6}_{1,1;j}} + x_{1,7}\widetilde{\Lambda^{1;5,6}_{2,1;j}}\\
\vdots\\
\widetilde{\Lambda^{1;5,6}_{j-1,1;j}} + x_{1,7}\widetilde{\Lambda^{1;5,6}_{j,1;j}}\\
\widetilde{\Lambda^{1;5,6}_{j,1;j}}\\
\widetilde{V^{3;5,8}_{j+1,1;j}}
\end{array}\right);\left(\begin{array}{c}
s_1\\
\vdots\\
s_{j-1}\\
s_j\\
(x_{1,3}-x_{1,7})(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4})
\end{array}\right)\right)_{R_2'} \{1-n\}\left\langle 1\right\rangle,
\end{aligned}
$$

where $R_2' = \mathbb{Q}[\mathcal{X}_{1,1}, \mathcal{X}_{j,2}, \mathcal{X}_{1,3}, \mathcal{X}_{j,4}, \mathcal{X}_{1,7}]/\widetilde{\Lambda^{1;5,6}_{j+1,1;j}}$. Using Corollary 2.11, there are polynomials $b_k \in R_2''$ $(k = 1, \cdots, j)$ and $b \in R_2'$ to give the following matrix factorization which is isomorphic to the above matrix factorization

$$
K\left(\left(\begin{array}{c}
b_1\\
\vdots\\
b_j\\
b
\end{array}\right);\left(\begin{array}{c}
s_1\\
\vdots\\
s_j\\
(x_{1,3}-x_{1,7})(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4})
\end{array}\right)\right)_{R_2'} \{1-n\}\left\langle 1\right\rangle.
$$

The partial matrix factorization $K(b;(x_{1,3}-x_{1,7})(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}))_{R_2'}$ is described as

$$
\begin{aligned}
& K\left(b;(x_{1,3}-x_{1,7})\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right)\right)_{R_2'}\\
\simeq\ & \left(\left(\begin{array}{r}
R_2''\\
(x_{1,3}-x_{1,7})R_2''\\
x_{1,7}(x_{1,3}-x_{1,7})R_2''\\
\vdots\\
x_{1,7}^{n-j-3}(x_{1,3}-x_{1,7})R_2''\\
x_{1,7}^{n-j-2}(x_{1,3}-x_{1,7})R_2''
\end{array}\right),\left(\begin{array}{r}
R_2''\{3-n\}\\
x_{1,7}R_2''\{3-n\}\\
x_{1,7}^2R_2''\{3-n\}\\
\vdots\\
x_{1,7}^{n-j-2}R_2''\{3-n\}\\
\beta R_2''\{3-n\}
\end{array}\right), g_0, g_1\right),
\end{aligned}
$$

where

$$\begin{aligned}
g_0 &= \begin{pmatrix} 0 & b(x_{1,3}-x_{1,7}) & & \\ \vdots & & \ddots & \\ 0 & & & b(x_{1,3}-x_{1,7}) \\ \frac{b}{\beta} & 0 & \cdots & 0 \end{pmatrix}, \\
g_1 &= \begin{pmatrix} 0 & \cdots & 0 & \beta(x_{1,3}-x_{1,7})(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}) \\ \sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4} & & & 0 \\ & \ddots & & \vdots \\ & & \sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4} & 0 \end{pmatrix}, \\
\beta &= \sum_{k=0}^{n-j-1} x_{1,7}^{n-j-1-k}\left(\sum_{l=0}^{k} x_{1,3}^{k-l}c_l\right), \\
R_2'' &= \mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{1,3},\mathcal{X}_{j,4}].
\end{aligned}$$

Remark that we have

$$\begin{aligned}
\beta(x_{1,3}-x_{1,7}) &= c_0x_{1,3}^{n-j}+c_1x_{1,3}^{n-j-1}+\cdots+c_{n-j}, \\
&\simeq \widetilde{\Lambda_{j+1,1;j}^{1;5,6}}\,|_{x_{1,7}\to x_{1,3}}\ .
\end{aligned}$$

Therefore the partial matrix factorization is isomorphic to

$$\begin{aligned}
&\bigoplus_{k=1}^{n-j-1} K\left(b(x_{1,3}-x_{1,7});\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right)_{R_2''}\{2k\} \\
&\qquad \bigoplus K\left(\frac{b}{\beta};\beta(x_{1,3}-x_{1,7})\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right)\right)_{R_2''}.
\end{aligned}$$

Then the total matrix factorization is isomorphic to

$$\bigoplus_{k=1}^{n-j-1} K\left(\begin{pmatrix} b_1 \\ \vdots \\ b_j \\ b(x_{1,3}-x_{1,7}) \end{pmatrix};\begin{pmatrix} s_1 \\ \vdots \\ s_j \\ \sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4} \end{pmatrix}\right)_{R_{2,}''}\{2k+1-n\}\langle 1\rangle \tag{12}$$

$$\bigoplus K\left(\begin{pmatrix} b_1 \\ \vdots \\ b_j \\ \frac{b}{\beta} \end{pmatrix};\begin{pmatrix} s_1 \\ \vdots \\ s_j \\ \widetilde{\Lambda_{j+1,1;j}^{1;5,6}}\,|_{x_{1,7}\to x_{1,3}}\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right) \end{pmatrix}\right)_{R_{2,}''}\{1-n\}\langle 1\rangle\,. \tag{13}$$

On the other hand, we have

$\mathcal{C}$



$$
\begin{aligned}
\mathcal{C}\left(\cdots\right)_n &= K\left(\begin{pmatrix}\Lambda^{1;9,2}_{1,1:j-1}\\ \vdots \\ \Lambda^{1;9,2}_{j,1:j-1}\end{pmatrix};\begin{pmatrix} x_{1,2}-X^{1;9}_{1,1;j-1}\\ \vdots\\ x_{j,2}-X^{1;9}_{j,1;j-1}\end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,1},\mathcal{X}_{j,2},\mathcal{X}_{j-1,9}]}\{1-j\}\\
&\quad\boxtimes K\left(\begin{pmatrix}V^{3;9,4}_{1,1:j-1}\\ \vdots \\ V^{3;9,4}_{j,1:j-1}\end{pmatrix};\begin{pmatrix} X^{3;9}_{1,1;j-1}-x_{1,4}\\ \vdots\\ X^{3;9}_{j,1;j-1}-x_{j,4}\end{pmatrix}\right)_{\mathbb{Q}[\mathcal{X}_{1,3},\mathcal{X}_{j,4},\mathcal{X}_{j-1,9}]}\\
&\simeq K\left(\begin{pmatrix}\widetilde{\Lambda^{1;9,2}_{1,1:j-1}}\\ \vdots \\ \widetilde{\Lambda^{1;9,2}_{j-2,1:j-1}}\\ \widetilde{\Lambda^{1;9,2}_{j-1,1:j-1}}\\ \widetilde{\Lambda^{1;9,2}_{j,1:j-1}}\end{pmatrix};\begin{pmatrix} s_1\\ \vdots\\ s_{j-1}\\ s_j+\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\\ -\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\end{pmatrix}\right)_{R_2''}\{1-j\}\\
&\simeq K\left(\begin{pmatrix}\widetilde{\Lambda^{1;9,2}_{1,1:j-1}}\\ \vdots \\ \widetilde{\Lambda^{1;9,2}_{j-1,1:j-1}}\\ \widetilde{\Lambda^{1;9,2}_{j,1:j-1}}-\widetilde{\Lambda^{1;9,2}_{j-1,1:j-1}}\end{pmatrix};\begin{pmatrix} s_1\\ \vdots\\ s_j\\ -\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\end{pmatrix}\right)_{R_2''}\{1-j\}.
\end{aligned}
$$

Thus using Theorem 2.10, the partial matrix factorization (12) is isomorphic to

$$\bigoplus_{k=1}^{n-j-1}\mathcal{C}\left(\cdots\right)_n\{2k+j-n\}\langle 1\rangle.$$



Moreover we have

$$
\begin{aligned}
&\widetilde{\Lambda^{1;5,6}_{j+1,1;j}}\,\Big|_{x_{1,7}\to x_{1,3}}\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right)\\
=\ &\left.\frac{F_{j+1}(X^{1;5}_{1,1;j},\cdots,X^{1;5}_{j,1;j},x_{j+1,6})-F_{j+1}(X^{1;5}_{1,1;j},\cdots,X^{1;5}_{j,1;j},X^{1;5}_{j+1,1;j})}{x_{j+1,6}-X^{1;5}_{j+1,1;j}}\right|_{x_{1,7}\to x_{1,3}}\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right).
\end{aligned}
$$

Since we have

$$\begin{aligned}
X^{1;5}_{k,1;j}\Big|_{x_{1,7}\to x_{1,3}} &= x_{1,1}x_{k-1,5}+x_{k,5}|_{x_{1,7}\to x_{1,3}}\\
&= x_{1,1}\left(\left(\sum_{l=0}^{k-2}(-1)^{k-2-l}x_{1,3}^{k-2-l}x_{l,4}\right)x_{1,7}+\sum_{l=0}^{k-1}(-1)^{k-1-l}x_{1,3}^{k-1-l}x_{l,4}\right)\\
&\quad +\left(\sum_{l=0}^{k-1}(-1)^{k-1-l}x_{1,3}^{k-1-l}x_{l,4}\right)x_{1,7}+\sum_{l=0}^{k}(-1)^{k-l}x_{1,3}^{k-l}x_{l,4}\Bigg|_{x_{1,7}\to x_{1,3}}\\
&= x_{1,1}x_{k-1,4}+x_{k,4}=X^{1;4}_{k,1;j},
\end{aligned}$$

$\widetilde{\Lambda^{1;5,6}_{j+1,1;j}}\,\big|_{x_{1,7}\to x_{1,3}}\ \left(\sum_{l=0}^{j}(-1)^{j+1-l}x_{1,3}^{j-l}x_{l,4}\right)$ equals

$$\begin{aligned}
&\frac{F_{j+1}(X^{1;4}_{1,1;j},\cdots,X^{1;4}_{j,1;j},x_{1,3}x_{j,2})-F_{j+1}(X^{1;4}_{1,1;j},\cdots,X^{1;4}_{j,1;j},X^{1;4}_{j+1,1;j})}{x_{j+1,6}-X^{1;4}_{j+1,1;j}}\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right)\\
\equiv\ &\frac{F_{j+1}(X^{1;4}_{1,1;j},\cdots,X^{1;4}_{j,1;j},x_{1,3}(\sum_{l=0}^{j}(-1)^{j+1-l}x_{1,3}^{j-l}X^{1;4}_{l,1;j}))-F_{j+1}(X^{1;4}_{1,1;j},\cdots,X^{1;4}_{j,1;j},X^{1;4}_{j+1,1;j})}{x_{1,3}(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}X^{1;4}_{l,1;j})-X^{1;4}_{j+1,1;j}}\\
&\qquad\times\left(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}x_{l,4}\right)\qquad\qquad\mod\ \langle s_j\rangle_{R_2''}\\
=\ &\frac{F_{j+1}(X^{1;4}_{1,1;j},\cdots,X^{1;4}_{j,1;j},x_{1,3}(\sum_{l=0}^{j}(-1)^{j-l}x_{1,3}^{j-l}X^{1;4}_{l,1;j}))-F_{j+1}(X^{1;4}_{1,1;j},\cdots,X^{1;4}_{j,1;j},X^{1;4}_{j+1,1;j})}{x_{1,3}-x_{1,1}}\\
=\ &\frac{F_1(x_{1,3})-F_1(x_{1,1})}{x_{1,3}-x_{1,1}}+\frac{F_j(u_1,\cdots,u_j)-F_j(x_{1,4},\cdots,x_{j,4})}{x_{1,3}-x_{1,1}},
\end{aligned}$$

where $u_k=\displaystyle\sum_{l=0}^{k}(-1)^{k-l}x_{1,3}^{k-l}X^{1;4}_{l,1;j}$.

The second term $\dfrac{F_j(u_1,u_2,\cdots,u_j)-F_j(x_{1,4},x_{2,4},\cdots,x_{j,4})}{x_{1,3}-x_{1,1}}$ equals

$$\begin{aligned}
&\frac{F_j(u_1,u_2,\cdots,u_j)-F_j(x_{1,4},u_2,\cdots,u_j)}{x_{1,3}-x_{1,1}}\\
&+\frac{F_j(x_{1,4},u_2,u_3,\cdots,u_j)-F_j(x_{1,4},x_{2,4},u_3,\cdots,u_j)}{x_{1,3}-x_{1,1}}\\
&+\cdots+\frac{F_j(x_{1,4},\cdots,x_{j-1,4},u_j)-F_j(x_{1,4},\cdots,x_{j-1,4},x_{j,4})}{x_{1,3}-x_{1,1}}.
\end{aligned}$$

Since $u_k - x_{k,4} = (x_{1,1} - x_{1,3})\left(\sum_{l=0}^{k-1}(-1)^{k-1-l}x_{1,3}^{k-1-l}x_{l,4}\right)$ and $s_k = x_{k,2} - u_k$, the above polynomial equals

$$
\begin{aligned}
&\frac{F_j(u_1,u_2,\cdots,u_j) - F_j(x_{1,4},u_2,\cdots,u_j)}{u_1 - x_{1,4}}\\
&+\frac{F_j(x_{1,4},u_2,u_3,\cdots,u_j) - F_j(x_{1,4},x_{2,4},u_3,\cdots,u_j)}{u_2 - x_{2,4}}\left(-\sum_{l=0}^{1}(-1)^{1-l}x_{1,3}^{1-l}x_{l,4}\right)\\
&+\cdots+\frac{F_j(x_{1,4},\cdots,x_{j-1,4},u_j) - F_j(x_{1,4},\cdots,x_{j-1,4},x_{j,4})}{u_j - x_{j,4}}\left(-\sum_{l=0}^{j-1}(-1)^{j-1-l}x_{1,3}^{j-1-l}x_{l,4}\right)\\
\equiv\ &\frac{F_j(x_{1,2},x_{2,2},\cdots,x_{j,2}) - F_j(x_{1,4},x_{2,2},\cdots,x_{j,2})}{x_{1,2} - x_{1,4}}\\
&+\frac{F_j(x_{1,4},x_{2,2},x_{3,2},\cdots,x_{j,2}) - F_j(x_{1,4},x_{2,4},x_{3,2},\cdots,x_{j,2})}{x_{2,2} - x_{2,4}}\left(-\sum_{l=0}^{1}(-1)^{1-l}x_{1,3}^{1-l}x_{l,4}\right)\\
&+\cdots+\frac{F_j(x_{1,4},\cdots,x_{j-1,4},x_{j,2}) - F_j(x_{1,4},\cdots,x_{j-1,4},x_{j,4})}{x_{j,2} - x_{j,4}}\left(-\sum_{l=0}^{j-1}(-1)^{j-1-l}x_{1,3}^{j-1-l}x_{l,4}\right)\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\mod \langle s_1,s_2,\cdots,s_j\rangle_{R_2''}\\
=\ &-L_{1,j}^{2;4} - L_{2,j}^{2;4}\left(\sum_{l=0}^{1}(-1)^{1-l}x_{1,3}^{1-l}x_{l,4}\right) - \cdots - L_{j,j}^{2;4}\left(\sum_{l=0}^{j-1}(-1)^{j-1-l}x_{1,3}^{j-1-l}x_{l,4}\right).
\end{aligned}
$$

Hence using Theorem 2.10 and Proposition 2.9, the matrix factorization (13) is isomorphic to

$$
\begin{aligned}
&K\left(\begin{pmatrix} * \\ \vdots \\ * \\ \frac{b}{\beta} \end{pmatrix};\begin{pmatrix} s_1 \\ \vdots \\ s_j \\ L_{1,1}^{3;1} - \sum_{k=1}^{j} L_{k,j}^{2;4}\left(\sum_{l=0}^{k-1}(-1)^{k-1-l}x_{1,3}^{k-1-l}x_{l,4}\right) \end{pmatrix}\right)_{R_2{}'_{,}{}'}\{1-n\}\langle 1\rangle\\
\simeq\ &K\left(\begin{pmatrix} L_{1,j}^{2;4} \\ L_{2,j}^{2;4} \\ \vdots \\ L_{j,j}^{2;4} \\ x_{1,3}-x_{1,1} \end{pmatrix};\begin{pmatrix} x_{1,2}-x_{1,4}+x_{1,3}-x_{1,1} \\ x_{2,2}-x_{2,4}+(x_{1,3}-x_{1,1})\left(\sum_{l=0}^{1}(-1)^{1-l}x_{1,3}^{1-l}x_{l,4}\right) \\ \vdots \\ x_{j,2}-x_{j,4}+(x_{1,3}-x_{1,1})\left(\sum_{l=0}^{j-1}(-1)^{j-1-l}x_{1,3}^{j-1-l}x_{l,4}\right) \\ L_{1,1}^{3;1} - \sum_{k=1}^{j} L_{k,j}^{2;4}\left(\sum_{l=0}^{k-1}(-1)^{k-1-l}x_{1,3}^{k-1-l}x_{l,4}\right) \end{pmatrix}\right)_{R_2{}'_{,}{}'}\{1-n\}\langle 1\rangle\\
\simeq\ &K\left(\begin{pmatrix} L_{1,j}^{2;4} \\ \vdots \\ L_{j,j}^{2;4} \\ x_{1,3}-x_{1,1} \end{pmatrix};\begin{pmatrix} x_{1,2}-x_{1,4} \\ \vdots \\ x_{j,2}-x_{j,4} \\ L_{1,1}^{3;1} \end{pmatrix}\right)_{R_2{}'_{,}{}'}\{1-n\}\langle 1\rangle
\end{aligned}
$$

$$
\simeq K\left(\begin{pmatrix} L_{1,j}^{2;4} \\ \vdots \\ L_{j,j}^{2;4} \\ L_{1,1}^{3;1} \end{pmatrix};\begin{pmatrix} x_{1,2}-x_{1,4} \\ \vdots \\ x_{j,2}-x_{j,4} \\ x_{1,3}-x_{1,1} \end{pmatrix}\right)_{R_2{}'_{,}{}'} \simeq \mathcal{C}'\left(\begin{tikzcd} \mathcal{X}_{1,1} \arrow[d, "1"'] & \mathcal{X}_{j,2} \\ \mathcal{X}_{1,3} & \mathcal{X}_{j,4} \arrow[u, "j"'] \end{tikzcd}\right)_n .
$$

□

Corollary 3.10.



Proof. We consider the following matrix factorization

(14)



Using Proposition 3.9 (2) and Proposition 3.8 (1), the matrix factorization is isomorphic to



(15)

On the other hand, using Proposition 3.7, Proposition 3.8 (1) and Proposition 3.9 (2), the matrix factorization (14) is isomorphic to

$\mathcal{C}\Bigg($



$\Bigg)_n$

$\simeq \bigoplus_{i=0}^{(j_1-j_2)(j_2-1)} \Bigg(\mathcal{C}\Bigg($



$\Bigg)_n \{-(j_1-j_2)(j_2-1)+2i\} \Bigg)^{p(i,j_1-1;j_2-1)}$

$\simeq \bigoplus_{i=0}^{(j_1-j_2)(j_2-1)} \Bigg(\mathcal{C}\Bigg($



$\Bigg)_n \{-(j_1-j_2)(j_2-1)+2i\} \Bigg)^{p(i,j_1-1;j_2-1)}$

(16) $\bigoplus_{i=0}^{(j_1-j_2)(j_2-1)} \Bigg(\Bigg(\bigoplus_{k=1}^{j_1-1} \mathcal{C}\Bigg($



$\Bigg)_n \{2k-j_1\} \Bigg) \{-(j_1-j_2)(j_2-1)+2i\} \Bigg)^{p(i,j_1-1;j_2-1)}$.

The $\mathbb{Z}$-grading shift of the matrix factorization (15) is derived from $[j_2-1]\begin{bmatrix} j_1 \\ j_2 \end{bmatrix}$ and one of the matrix factorization (16) is also derived from $[j_1-1]\begin{bmatrix} j_1-1 \\ j_2-1 \end{bmatrix}$. Since $[j_1-1]\begin{bmatrix} j_1-1 \\ j_2-1 \end{bmatrix} - [j_2-1]\begin{bmatrix} j_1 \\ j_2 \end{bmatrix} = \begin{bmatrix} j_1-1 \\ j_2 \end{bmatrix}$, we obtain Corollary 3.10. ☐

Conjecture 3.11 (For Strategy (S1))**.** *There are equivalences in the category* HMF^{gr} *corresponding to the remaining MOY relations.*

For crossings with coloring $(1,2)$ and $(2,1)$, the author categorified MOY polynomial using the category Kom(HMF) [11].

REFERENCES

[1] D. Bar-Natan, Khovanov's homology for tangles and cobordisms, Geom. Topol.**9** (2005), 1443–1499 .
[2] D. Gepner, Fusion rings and geometry, Comm. Math. Phys. 141 (1991), no. 2, 381–411.
[3] B. Gornik, Note on Khovanov link cohomology, arXiv:math.QA/0402266.
[4] M. Khovanov, A categorification of the Jones polynomial, Duke Math. J. **101** (2000), no. 3, 359-426.
[5] M. Khovanov, L. Rozansky, Matrix factorizations and link homology, arXiv:math.QA/0401268.
[6] M. Khovanov, L. Rozansky, Matrix factorizations and link homology II, arXiv:math.QA/0505056.
[7] M. Khovanov, L. Rozansky, Virtual crossings, convolutions and a categorification of the SO(2N) Kauffman polynomial, arXiv:math.QA/0701333

[8] H. Murakami, T. Ohtsuki, S. Yamada, Homfly polynomial via an invariant of colored plane graphs, Enseign. Math. (2) **44** (1998), no. 3-4, 325–360.
[9] J. Rasmussen, Some differentials on Khovanov-Rozansky homology, arXiv:math.GT/0607544.
[10] Y. Yonezawa, Matrix factorizations and double line in $\mathfrak{sl}_n$ quantum link invariant, arXiv:math.GT/0703779.
[11] Y. Yonezawa, A categorification of MOY polynomial with coloring 1 and 2.
[12] H. Wu, On the quantum filtration of the Khovanov-Rozansky cohomology, arXiv:math.GT/0612406.
[13] H. Wu, Matrix factorizations and colored MOY graphs, arXiv:0803.2071.

Graduate School of Mathematics, Nagoya University, 464-8602 Furocho, Chikusaku, Nagoya, Japan
E-mail address: `yasuyoshi.yonezawa@math.nagoya-u.ac.jp`